arXiv:math/9807177v1 [math.LO] 15 Jul 1998

PCF AND INFINITE FREE SUBSETS IN AN ALGEBRA
SH513

Saharon Shelah

Institute of Mathematics
The Hebrew University
Jerusalem, Israel

Rutgers University
Mathematics Department
New Brunswick, NJ USA

I would like to thank Alice Leonhardt for the beautiful typing.
Research partially supported by the Israel Science Foundation
First Typed at Rutgers - 98/July/16
Done - 6-7/93
Previous version - 98/Mar/19

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

Anotated Content

§1 Other variants of "G.C.H. holds almost always"

[We give another way to prove that for every $\lambda \geq \beth_\omega$ for every large enough regular $\kappa < \beth_\omega$ we have $\lambda^{[\kappa]} = \lambda$, dealing with sufficient conditions for replacing $\beth_\omega$ by $\aleph_\omega$.]

§2 Large pcf($\mathfrak{a}$) implies the existence of free sets

[A nice example of the implication stated in the title is that if $\mathrm{pp}(\aleph_\omega) > \aleph_{\omega_1}$ then for every algebra M of cardinality $\aleph_\omega$ with countably many functions (or just $< \aleph_\omega$ many), for some $a_n \in M$ (for $n < \omega$) we have $a_n \notin c\ell_M(\{a_\ell : \ell \neq n, \ell < \omega\})$. Generally if $\mathrm{pcf}(\mathfrak{a})$ is not just of cardinality $> |\mathfrak{a}|$, but $\langle J_{<\theta}[\mathfrak{a}] : \theta \in \mathrm{pcf}(\mathfrak{a})\rangle$ has large rank (as defined below) than a relevant instance of IND connected to $\sup(\mathfrak{a})$ holds.]

§3 Existence of free subsets implies restrictions on pcf

[We have results of forms complementary to those of §2 (though not close enough). So if $\mathrm{IND}(\mu, \sigma)$ (in every algebra with universe λ and $\leq \sigma$ functions there is an infinite independent subset) then for no distinct regular $\lambda_i \in \mathrm{Reg} \backslash \mu^+$ (for $i < \kappa$) does $\prod\limits_{i<\kappa} \lambda_i / [\kappa]^{\leq \sigma}$ have true cofinality. We also look at $\mathrm{IND}(\langle J^{\mathrm{bd}}_{\kappa_n} : n < \omega\rangle), J_n$ an ideal on κ_n (we ask for $\bar{\alpha} \in \prod\limits_{n<\omega} \kappa_n$ such that $n < \omega \Rightarrow \alpha_n > \sup c\ell_\mu(\{\alpha_\ell : \ell \in (n, \omega)\})$ and more general version, and from assumptions as in §2 get results even for the non stationary ideal.]

§4 Sticks and Boolean Algebras

[We deal with some other measurements of $[\lambda]^{\geq \theta}$ and give an application by a construction of a Boolean Algebra.]

§5 More on independence

§6 Odds and ends

[In 6.1 we deal with a replacement for Δ-system lemma. We have $> 2^\kappa$ sequences of length κ. In 6.3 we look at how we can divide $F \subseteq \Pi_\alpha$ to few bounded sets. In 6.4 we relook at the characterization of a property of [GHS], generalizing the questions somewhat. We then deal with freeness properties for $F \subseteq {}^\delta\mathrm{Ord}$ (modulo an ideal) and we give a correct version of [Sh:g, Ch.IX,3.5] on characterizing $\mathrm{cov}(\lambda, \lambda, \theta, \sigma)$ when $\sigma > \aleph_0$ concerning the obtainment of the pp version. We shall continue in [Sh 589].]

§1 Other variants of “G.C.H. holds almost always”

We essentially redo the proof of [Sh 460], §2 in another more general way.

1.1 Notation. 1) $\mathfrak{F}_\kappa(A)$ is the family of κ-complete filters $\mathfrak{D}$ on $\mathscr{P}(A)$ (so $\mathfrak{D} \subseteq \mathscr{P}(\mathscr{P}(A))$ (so the points are subsets of A, and the members of $\mathfrak{D}$ (which are $\subseteq \mathscr{P}(A)$) which we shall be most interested in are ideal and their compliments.
2) We say $\mathfrak{D} \in \mathfrak{F}_\kappa(A)$ has σ-complete character if for any $Y \subseteq \mathscr{P}(A)$ we have: $Y \in \mathfrak{D}$ iff $\mathrm{fil}_\sigma(Y) \in \mathfrak{D}$ where $\mathrm{fil}_\sigma(Y)$ is the σ-complete ideal on A generated by Y.
For an ideal I on $X, I^+ = \mathscr{P}(X) \backslash I$, similarly for a filter.

1.2 Definition. 1) For $\mathfrak{D} \in \mathfrak{F}_\kappa(A)$, cardinals $\mu < \lambda$ and σ such that $|A| < \mu < \lambda$, we say that λ is $(\mathfrak{D}, \mu, \sigma)$-inaccessible when: if $\mathfrak{a}_t \subseteq (\mu, \lambda) \cap$ Reg for $t \in A, |\mathfrak{a}_t| < \sigma$ then $\{B \subseteq A : \mathrm{pcf}_{\sigma\text{-complete}}(\cup\{\mathfrak{a}_t : t \in B\}) \subseteq \lambda\} \in \mathfrak{D}$.
2) If we omit μ we mean: for $\mu = (|A| + \sigma)^+$.

1.3 Theorem. *Suppose $\langle \kappa_n : n < \omega \rangle$ is a strictly increasing sequence of regular cardinals $> \aleph_2$. Stipulate $\kappa_{-1} = \aleph_1$ and assume $\mathfrak{D}_n \in \mathfrak{F}_{\kappa_{n-1}}(\kappa_n)$ for $n < \omega$ and $\kappa = \sum\limits_{n<\omega} \kappa_n$ satisfies:*

> $\otimes$ *if $n < \omega, \aleph_0 < \theta =$ cf$(\theta) < \kappa_n, h : \kappa_{n=1} \to \theta$ and*
> *$Y \in \mathfrak{D}^+_{n+1}$ (so $Y \subseteq \mathscr{P}(\kappa_{n+1})$) then for some $\zeta < \theta$ we have*
> $(*)_{y,\zeta}$ $\{B \in Y : \text{sup Rang}(h \restriction B) < \zeta\} \in \mathfrak{D}^+_{n+1}$.

If $\lambda > \kappa$ then for every $n < \omega$ large enough, λ is $(\mathfrak{D}_n, \kappa, \aleph_1)$-inaccessible.

1.4 Remark. 1) We can replace $\omega, \aleph_1$ by θ, θ^+ or $< \theta, \theta$ (when θ is regular uncountable) respectively (so $\kappa = \sum i < \theta \kappa_i$, etc.) (why? repeat the proof or force by Levy$(\aleph_0, < \sigma)$). Of course, we can replace $\mathfrak{F}(\kappa_n)$ by $\mathfrak{F}(A)$ if $|A| = \kappa_n$.
2) Note that the set defined in 1.2(2) is always an ideal on A.

Proof. We prove this by induction on λ. If $\lambda = \kappa^+$ this is empty (as $(\kappa, \lambda) = \emptyset$). Also if $\lambda < \kappa^{+\omega_1}$ this is trivial, as $\bigcup\limits_{t \in A} \mathfrak{a}_t$ is countable $(\subseteq \{\kappa^{+(\alpha+1)} : \alpha < \omega_1\} \cap \lambda)$ hence $\mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup\limits_{t \in A} \mathfrak{a}_t) = \bigcup\limits_{t \in A} \mathfrak{a}_t \subseteq \{\kappa^{+(\alpha+1)} : \alpha < \omega_1\} \cap \lambda$. Also if this holds for λ it holds for λ^+ because $\mathrm{pcf}(\mathfrak{a} \cup \{\lambda\}) \subseteq \mathrm{pcf}(\mathfrak{a}) \cup \{\lambda\}$. So we can assume that λ is a limit cardinal. If the conclusion fails then for some infinite $W \subseteq \omega$, for $n \in W$ we have a sequence $\langle \mathfrak{a}^n_\alpha : \alpha < \kappa_n \rangle$ (where $\mathfrak{a}^n_\alpha \subseteq (\kappa, \lambda) \cap$ Reg) which is a counterexample, i.e. $Y_n =: \{B \subseteq \kappa_n : \lambda \not\supseteq \mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup\limits_{\alpha \in B} \mathfrak{a}^n_\alpha)\} \in \mathfrak{D}^+_n$. If cf$(\lambda) > \kappa$ then $\cup\{\mathfrak{a}^n_\alpha : n < \omega, \alpha < \kappa_n\}$ is a subset of λ of cardinality $\leq \kappa$, hence is bounded by some $\lambda' < \lambda$, so apply the induction hypothesis on λ'. If $\aleph_0 <$ cf$(\lambda) < \kappa$ let $\lambda = \sum\{\lambda_\zeta : \zeta <$ cf$(\lambda)\}$, $\bigwedge\limits_{\zeta < \xi} \lambda_\zeta < \lambda_\xi < \lambda$. Now as cf$(\kappa) = \aleph_0$ and $\kappa = \sum\limits_{n<\omega} \kappa_n$, for some $n(*) < \omega$ we have $\mathrm{cf}(\lambda) < \kappa_{n(*)}$. For every $n \in W\backslash(n(*)+2)$, we define a function $h_n : \kappa_n \to \mathrm{cf}(\lambda)$ by $h_n(\alpha) = \mathrm{Min}\{\zeta < \kappa_{n(*)} : \mathfrak{a}_\alpha \subseteq \lambda_\zeta\}$. Hence by the assumption $\otimes$, as $\mathrm{cf}(\lambda) < \kappa_{n(*)} < \kappa_{n-1}$, for some $\zeta_n < \mathrm{cf}(\lambda)$ we have $\{B \subseteq \kappa_n : \bigwedge_{\alpha\in B} \mathfrak{a}_\alpha \subseteq \lambda_{\zeta_n}\} \in \mathfrak{D}^+_n$. Now we can contradict the induction hypothesis for $\lambda' = \sup\{\lambda_{\zeta_n} : n \in W\backslash(n(*)+2)\}$. We are left with the case $\mathrm{cf}(\lambda) = \aleph_0$ so let $\lambda = \sum_{n<\omega} \lambda_n, \lambda_0 = \kappa^+, \lambda_n < \lambda_{n+1}$. Note that as our $\mathfrak{D}_m$ is κ_{m-1}-complete (see 1.1) hence is $\aleph_1$-complete, without loss of generality for some $\langle \lambda_k : k < \omega\rangle$ strictly increasing and $\lambda = \sum_{k<\omega} \lambda_k, \lambda_0 = \kappa^+$ define for each $n, k < \omega, Y^k_n = \{B \subseteq \kappa_n : \lambda \not\supseteq \mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup_{\alpha\in B} \mathfrak{a}^n_i \cap [\lambda_k, \lambda_{k+1})\}$. So $Y_n = \bigcup_{k<\omega} Y^k_n$, but $Y_n \in \mathfrak{D}^+_n$, and $\mathfrak{D}^+_n$ is $\aleph_1$- complete hence for some $k_n < \omega, Y^{k_n}_n \in \mathfrak{D}^+_n$, so possibly shrinking W we get $\langle k_n : n \in W\rangle$ is constant or strictly increasing, the former contradicts the induction hypothesis on $\lambda_{k_{\mathrm{Min}(W)+1}}$. Now renaming the λ_k's we get $k_n = n$ and we can replace $\mathfrak{a}^n_\alpha$ by $a^n_\alpha \cap [\lambda_n, \lambda_{n+1})$. So without loss of generality $\mathrm{Min}(W) > 4$ and for $n \in W$ we have $\bigcup_\alpha \mathfrak{a}^n_\alpha \subseteq [\lambda_n, \lambda_{n+1})$ and $\lambda_n < \lambda$, of course.

Let $n(*) = \min(W)$. We try to define by induction on $k < \omega, \langle \theta_t : t \in \omega_k\rangle, w_k = \bigcup_{i<\kappa_{n(*)}} w_{k,i}, J_k$ and h_{k-1} if $k > 0$ such that:

(a) $\theta_t \in \mathrm{Reg} \cap \lambda\backslash\kappa$ for $t \in w_k$

(b) $w_k = \bigcup_{i<\kappa_{n(*)}} w_{k,i}$ is disjoint to $\bigcup_{\ell<k} w_\ell$

(c) $\langle w_{k,i} : i < \kappa_{n(*)}\rangle$ is a sequence of pairwise disjoint, countable sets sn

(d) $w_{0,i} = \{i\} \times \mathfrak{a}^{n(*)}_i$ and $\theta_{(i,\tau)} = \tau$ for $\tau \in \mathfrak{a}^{n(*)}_i$

(e) h_k is a function from w_{k+1} to w_k mapping $w_{k+1,i}$ into $w_{k,i}$

(f) $J_k = \{w \subseteq w_k : \lambda \supseteq \mathrm{pcf}_{\aleph_2\text{-complete}}(\{\theta_t : t \in w\})\}$ is a proper ideal

(g) if $w \in J^+_k$ then $\{t \in w_{k+1} : h_k(t) \in w\} \in J^+_{k+1}$

(h) $t \in w_{k+1} \Rightarrow \theta_t < \theta_{h_k(t)}$.

During the induction, h_k is defined in the k-th step.

If we succeed, we shortly get a contradiction (by observation [Sh 460, 2.2]). For $k = 0$ define $w_{0,i}, \theta_\tau$ for $\tau \in w_0 = \bigcup_{i<\kappa_{n(*)}} w_{0,i}$ by clause (d) and the clause (f) holds as otherwise $\{\theta_t : t \in w_0\}$ can be represented as $\bigcup_{\varepsilon<\omega_1} \mathfrak{b}_\varepsilon$ with $\max \mathrm{pcf}(\mathfrak{b}_\varepsilon) < \lambda$, let $h : \kappa_{n(*)} \to \omega_1$ be $h(i) = \sup\{\min\{\varepsilon : \tau \in \mathfrak{b}_\varepsilon\} : \tau \in \mathfrak{a}^{n(*)}_i\}$ and apply $\otimes$ from the hypothesis to get $Y \subseteq Y_{n(*)}$ and $\zeta < \omega_1$ such that $Y \in \mathfrak{D}^+_{n(*)}$ and $\{B \in Y : \sup \mathrm{Rang}(h \restriction B) < \zeta\} \in \mathfrak{D}^+_{n(*)}$, but $B \in Y$ implies $\lambda \not\supseteq \mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup_{\alpha\in B} \mathfrak{a}^{n(*)}_\alpha)$ because $B \in Y_{n(*)}$ and

$$\mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup_{\alpha\in B} \mathfrak{a}^{n(*)}_\alpha) \subseteq \mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup_{\varepsilon<\zeta} \mathfrak{b}_\varepsilon) \subseteq \bigcup_{\varepsilon<\zeta} \mathrm{pcf}_{\aleph_1\text{-complete}}(\mathfrak{b}_\varepsilon) \subseteq \lambda;$$

contradiction. So assume $w_k, \langle w_{k,i} : i < \kappa_{n(*)}\rangle, J_k$ are as required and we shall define $w_{k+1}, \langle w_{k+1,i} : i < \kappa_{n(*)}\rangle, J_{k+1}, h_k$.
Now for any $t \in w_k$ by the induction hypothesis for some $g(t) < \omega$ we have

$(*)_1$ if $m \in [g(t), \omega)$ and $\mathfrak{b}_i \subseteq$ Reg $\cap\, \theta_t \backslash \kappa$ is countable for $i < \kappa_m$ then

$$\{B \subseteq \kappa_m : \mathrm{pcf}_{\aleph_1\text{-complete}}(\cup\{\mathfrak{b}_i : i \in B\}) \subseteq \theta_t\} \in \mathfrak{D}_m$$

$(*)_2$ $g(t) > n(*) + 1$.

Let $u_{k,m} = \{t \in w_k : g(t) = m \text{ and } \theta_t > \kappa^+\}$. We shall prove

$\boxtimes_{k,m}$ if $u = u_{k,m} \notin J_k$, then we can find $\langle \mathfrak{c}_t : t \in u_{k,m}\rangle, \mathfrak{c}_t \subseteq$ Reg $\cap \theta_t \backslash k$ countable such that:
$u \subseteq u_{k,m}, u \in J^+_\kappa$ implies $\mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup\limits_{t \in u} \mathfrak{c}_t) \nsubseteq \lambda$.

As J_k is $\aleph_1$-complete (by its definition; even more) this suffices for carrying the induction.
[Why? Let $\langle \mathfrak{c}^m_t : t \in u_{k,m}\rangle$ for $m < \omega$ such that $u_{k,m} \notin J_k$ be as above, let

$$w_{k+1,i} = \bigcup\{\{t\} \times \mathfrak{c}^m_t : \text{for some } m, u_{k,m} \notin J_k \text{ and } t \in u_{k,m} \cap w_{k,i}\},$$

$\theta_{(t,\tau)} = \tau, w_{k+1} = \bigcup\limits_{i<k_{n(*)}} w_{k+1,i}$ and we define the function $h_{k+1} : w_{k+1} \rightarrow w_k$ by $h_{k+1}((t,\sigma)) = t$ (note: every t belongs to at most one $u_{k,m}\, (m < \omega)$ and $w_k \backslash \cup \{u_{k,m} : u_{k,m} \notin J_k\} = \emptyset \bmod J_k$.]

Proof of $\boxtimes_{k,m}$. For each $\tau \in \mathfrak{a}_m$, we apply [Sh:g, Ch.I,1.6] or [Sh:g, Ch.IX,4.1] on $\langle \theta_t : t \in u_{k,m}\rangle, J = J_k \restriction u_{k,m}$ and τ (possible as $|u_{k,m}| < \kappa <$ $\min\{\theta_t : t \in u_{k,m}\}$), each $\theta_t (t \in u_{k,m})$ regular and $\prod\limits_{t \in u_{k,m}} \theta_t / J$ is λ^+-directed, $\tau < \lambda^+$ (the cases $\tau < \theta_t$ can be ignored for several reasons, e.g. $\bigcup\limits_\alpha \mathfrak{a}^m_\alpha \subseteq [\lambda_m, \lambda_{m+1})$). So we can find $\langle \theta_{t,\tau} : \tau \in \mathfrak{a}_m\rangle$ such that:

(α) $\theta_{t,\tau}$ is regular $\kappa^+ \le \theta_{t,\tau} < \theta_t$

(β) $\prod\limits_{t \in u_{k,m}} \theta_{t,\tau}/J_k$ has true cofinality τ

(note that $t \in u_{k,m} \Rightarrow \kappa^+ < \theta_t$, so we can assume $\theta_{t,\tau} \ge \kappa^+$).

Now for each $t \in u_{k,m}, \theta_{t,\tau} \in$ Reg $\cap \theta_t \backslash \kappa$ for $\tau \in \mathfrak{a}_m$, but $g(t) = m$ (as $t \in u_{k,m}$), hence by the definition of g, letting $\mathfrak{a}^{m,t}_i = \{\theta_{t,\tau} : \tau \in \mathfrak{a}^m_i\}$, for $i < \kappa_m$, we have

$$\Gamma^m_t =: \{B \subseteq \kappa_m : \theta_t \supseteq \mathrm{pcf}_{\aleph_1\text{-complete}}(\bigcup_{i \in B} \mathfrak{a}^{m,t}_i)\} \in \mathfrak{D}_m.$$

But $\mathfrak{D}_m$ is $\kappa_{n(*)+1}$-complete (as $m = g(t) > n(*) + 1$) and $|u_{k,m}| \le \kappa_{n(*)} < \kappa_{n(*)+1}$ hence $\Gamma^* = \bigcap\limits_{t \in u_{k,m}} \Gamma^m_t \in \mathfrak{D}_m$. On the other hand (by the choice of $\mathfrak{a}_m$)

$$\Gamma_m =: \{B \subseteq \kappa_m : \lambda \supseteq \text{ pcf}_{\aleph_1\text{-complete}}(\bigcup_{i \in B} \mathfrak{a}^m_t)\} \notin \mathfrak{D}_m.$$

So there is $B \in \Gamma_m \cap \Gamma^* = \Gamma_m \cap \bigcap_{t \in u_{k,m}} \Gamma^m_t$.

Let

$$\mathfrak{a}^* = \{\theta_t : t \in u_{k,m}\} \cup \{\theta_{t,\tau} : t \in u_{k,m}, \tau \in \mathfrak{a}_m\} \cup \mathfrak{a}_m,$$

and for simplicity assume $\lambda \cap \text{ pcf}(\mathfrak{a}^*)| < \text{ min}(\mathfrak{a}^*)$ hence there is a smooth close generating sequence $\langle \mathfrak{b}_\sigma[\mathfrak{a}^*] : \sigma \in \lambda \cap \text{ pcf}(\mathfrak{a}^*)\rangle$ (see e.g. [Sh 430, 6.7], if not use [Sh 430], 6.7 - **?**). Clearly $B \neq \emptyset$. For each $t \in u_{k,m}$ we know $B \in \Gamma^m_t$ hence

−> scite{6.7F} undefined

$\theta_t \supseteq \text{ pcf}_{\aleph_1\text{-complete}}(\bigcup_{i \in B} \mathfrak{a}^{m,t}_i)$. So we can find countable $\mathfrak{c}_t \subseteq \theta_t \cap \text{ pcf}(\bigcup_{i \in B} \mathfrak{a}^{m,t}_i)$ such that

$$\bigcup_{i \in B} \mathfrak{a}^{m,t}_i \subseteq \bigcup_{\sigma \in \mathfrak{c}_t} \mathfrak{b}_\sigma[\mathfrak{a}^*].$$

The pcf calculus verifies clause (g) (as in [Sh 460, §2]). $\square_{1.3}$

It is now natural to look for suitable filters $\mathfrak{D}$, the simplest ones are.

1.5 Definition. For $\sigma < \theta < \kappa$ and $\mu \leq \kappa$ (always σ, θ regular) let $\mathfrak{D} = \mathfrak{D}_{\sigma,\theta,\kappa,\mu}$ be the following filter on $\mathscr{P}(\kappa) : Y \in \mathfrak{D}$ iff there are functions $f_\alpha : \kappa \to \theta_\alpha$ for $\alpha < \chi(*) < \theta$ where $\theta_\alpha \in [\sigma, \theta) \cap$ Reg such that $Y \supseteq \{a \subseteq \kappa : |a| \geq \mu$ and for every $\alpha < \theta$ for some $\zeta < \theta_\alpha$ we have $\text{Rang}(f_\alpha \restriction a) \subseteq \zeta\}$. If $\mu = \theta$ we may omit it.

1.6 Observation. 1) If $\sigma < \theta < \kappa_1 \leq \kappa_2$ and $\emptyset \in \mathfrak{D}_{\sigma,\theta,\kappa_1,\mu}$ then $\emptyset \notin \mathfrak{D}_{\sigma,\theta,\kappa_2,\mu}$.
2) $\mathfrak{D}_{\sigma,\theta,\kappa}$ is a θ-complete filter.
3) If $2^{<\theta} < \kappa$ then $\emptyset \notin \mathfrak{D}_{\sigma,\theta,\kappa}$ and this is preserved by σ-c.c. forcing.

Proof. Straight.

1.7 Conclusion. Let μ be a limit singular cardinal of cofinality $< \sigma < \mu$ and:

⊗ for every $\theta \in (\sigma, \mu)$ for some $\kappa \in (\theta, \mu)$ we have: $\emptyset \notin \mathfrak{D}_{\sigma,\theta,\kappa}$.

Then for every $\lambda > \mu$, for some $\theta = \theta_\mu \in (\sigma, \mu)$ for every $\kappa \in (\theta, \mu)$ we have

(∗) if $\lambda_i \in (\mu, \lambda) \cap$ Reg for $i < \kappa$ then

$$\{a \subseteq \kappa : \text{pcf}_{\sigma\text{-complete}}\{\lambda_i : i \in a\} \subseteq \lambda\} \in \mathfrak{D}_{\sigma,\theta,\kappa}.$$

Proof. Assume λ is a counterexample. Without loss of generalityσ is regular, choose by induction on $\zeta < \sigma, \kappa_\zeta \in (\sigma, \mu) \cap$ Reg as follows

$$\kappa_0 \in (\sigma, \mu) \cap \text{ Reg arbitrary;}$$

$$\kappa_\zeta \in (\bigcup_{\epsilon<\zeta} \kappa^+_\epsilon, \mu) \cap \text{ Reg is minimal } \kappa \text{ which is a counterexample to } \otimes \text{ for}$$

$$\theta = \bigcup_{\epsilon<\zeta} \kappa^+_\epsilon \text{ (in particular } \emptyset \notin \mathfrak{D}_{\sigma, \cup\{\kappa^+_\epsilon : \epsilon<\zeta\}, \kappa_\zeta}, \text{ so } \kappa_\zeta < \mu).$$

Let $\kappa = \bigcup\limits_{\zeta<\sigma} \kappa_\zeta$ and apply 1.3 for $\mathfrak{D}_{\sigma, \bigcup\limits_{\epsilon<\zeta} \kappa^+_\epsilon, \kappa_\zeta}$ from Definition 1.5, more exactly the variant with σ instead $\aleph_1$ (see 1.4). $\square_{1.7}$

1.8 Remark. 1) In the proof of 1.3 we can change the universe during the proof, so weaken the demand $\otimes$.
2) The problematic example is: $T \subseteq {}^{\omega_1}\omega$ a Kurepa tree, say $T \cap {}^\alpha 2 = \{\gamma_\alpha(n) : m < \omega\}, \eta_j \in \lim_{\omega_1}(T)$ for $j < j^*$ and $\{a \subseteq \omega_1 \cup j^* : \delta =: a \cap \omega_1$ a limit ordinal and for every $n < \omega$ for some $j \in j^* \cap a$ we have $\gamma_\alpha(n) = \eta_j(\alpha)\} \in \mathscr{D}_{<\aleph_1}(\omega_1 \cup j^*)$.
3) We can replace in 1.7 above $\text{cf}(\mu) < \sigma$ by $\text{cf}(\mu) \neq \sigma$. We can replace $\emptyset \notin \mathfrak{D}_{\sigma,\theta,\kappa}$ by $\emptyset \in \mathfrak{D}_{\sigma,\theta,\kappa;\Upsilon}$ for any fix $\Upsilon \in [\sigma, \mu)$. Note that the case $\Upsilon < \sigma$ is not interesting.
4) Note that the meaning of $\emptyset \in \mathfrak{D}_{\sigma,\theta,\kappa;\Upsilon}$ is that there are $\alpha(*) < \theta$ and functions $f_\alpha : \kappa \to \theta_\alpha$ where $\theta_\alpha \in$ Reg $\cap [\sigma, \theta)$ such that for no $u \in [\kappa]^\Upsilon$ do we have $\alpha < \alpha(*) \Rightarrow$ sup Rang$(f_\alpha \restriction u) < \theta_\alpha$. Recall if Υ is omitted it means $\Upsilon = \theta$.

1.9 Definition. Assume $\mathbf{J} \subseteq$ Id(κ) (= the family of ideals on κ).
1) We say (λ, μ) is **J**-inaccessible if $\kappa \leq \mu < \lambda$ and there are no $\lambda_i \in$ Reg $\cap (\mu, \lambda)$ for $i < \kappa$ and $J \in \mathbf{J}$ such that $\prod\limits_{i<\kappa} \lambda_i / \mathbf{J}$ is λ-directed (equivalently for some such λ_i's, $\prod\limits_{i<\kappa} \lambda_i / J$ has true cofinality and it is $\geq \lambda$).
2) $(\lambda, *)$ means (λ, μ) for some $\mu \in [\kappa, \lambda), \lambda$ means (λ, κ).
3) **J** is σ-indecomposable when: if $J \in \mathbf{J}$ and $h : \text{Dom}(J) \to \sigma$ then for some $\zeta < \sigma$ and $I \in \mathbf{J}$ we have $J \restriction h^{-1}\{\varepsilon : \varepsilon < \zeta\} \leq^* I$ (see below).
4) For ideals I_ℓ, on $A_\ell\,(\ell = 0, 1) I_0 \leq^* I_1$ if there is $B_0 \in I^+_0$ and $B_1 \in I^+_1$ and one-to-one function g from B_0 into B_1 such that

$$Y \cap B_0 \in I_0 \Rightarrow g''(Y \cap B_0) \in J_1$$

5) **J** is $[\sigma, \kappa)$-indecomposable if it is θ-indecomposable for every $\theta \in [\sigma, \kappa) \cap$ Reg.

§2 LARGE PCF($\mathfrak{a}$) IMPLIES THE EXISTENCE OF FREE SETS

2.1 Definition. 1) Let $\bar{A} = \langle A_\alpha : \alpha < \alpha^* \rangle$ be a sequence of subsets of κ, no A_α in the ideal generated by $\{A_\beta : \beta < \alpha\}$. We define functions $\mathrm{rk} = \mathrm{rk}_{\bar{A}}$, $\mathrm{rk}' = \mathrm{rk}'_{\bar{A}}$ from $\mathscr{P}(\kappa)$ to the ordinals by:

> $\mathrm{rk}(A) \geq \zeta$ iff for every $\xi < \zeta$ for some $\alpha, A \neq A \cap A_\alpha$ and $\mathrm{rk}(A \cap A_\alpha) \geq \zeta$
>
> $\mathrm{rk}'(A) \geq \zeta$ iff for every $\xi < \zeta$ for some α, we have $\mathrm{rk}'(A \cap A_\alpha) \geq \xi$ and $A \backslash A_\alpha, A \cap A_\alpha$ are not in $\mathrm{id}_{\bar{A} \restriction \alpha}$ = the ideal generated by $\{A_\beta : \beta < \alpha\}$.

2) Let $\bar{J} = \langle (J_\alpha, J'_\alpha) : \alpha < \alpha^* \rangle$ be a sequence of pair of ideals on κ such that $[\alpha < \beta \Rightarrow J_\alpha \subseteq J'_\alpha \subseteq J_\beta \subseteq J'_\beta]$ and for some $A_\alpha \in J^+_\alpha, J'_\alpha = J_\alpha + A_\alpha$ we define $\mathrm{rk}'_{\bar{J}}(A)$ for $A \subseteq \kappa$ by:

> $\mathrm{rk}'_{\bar{J}}(A) \geq \zeta$ iff for every $\xi < \zeta$ for some $\alpha < \alpha^*$ we have: $\mathrm{rk}'_{\bar{J}}(A \cap A_\alpha) \geq \xi$ and $A \backslash A_\alpha, A \cap A_\alpha$ are not in J_α.

3) We identify $\bar{A}$ with $\langle (\mathrm{id}_{\bar{A} \restriction \alpha}, \mathrm{id}_{\bar{A} \restriction (\alpha+1)}) : \alpha < \alpha^* \rangle$ (see 2.2(3) below). If $\bar{J} = \langle (J_\alpha, J_{\alpha+1}) : \alpha < \alpha^* \rangle$ is as required in (2) we may write $\langle J_\alpha : \alpha < \alpha^* \rangle$ instead $\bar{J}$. We can replace κ by any other set. We may write $\mathrm{rk}^{(\prime)}(A, \bar{A})$ or $\mathrm{rk}'(A, \bar{J})$ instead $\mathrm{rk}^{(\prime)}_{\bar{A}}(A)$ or $\mathrm{rk}'_{\bar{J}}(A)$.

2.2 Claim. *1) $rk_{\bar{A}}$,$rk'_{\bar{A}}$ are well defined (values: ordinals or ∞) and nondecreasing in A (under $\subseteq$).*
2) $rk_{\bar{A}}(A) \geq rk'_{\bar{A}}(A)$.
3) $rk'_{\bar{A}}$ depend just on $\langle id_{\bar{A} \restriction \alpha} : \alpha \leq \alpha^* \rangle$ and for $A \subseteq \kappa$, we have $rk'_{\bar{A}}(A) = rk'_{\bar{J}}(A)$ where $\bar{J} = \langle (id_{\bar{A} \restriction \alpha}, id_{\bar{A} \restriction (\alpha+1)}) : \alpha \leq \alpha^* \rangle$ (so we may write $rk'_{\langle J_\alpha : \alpha \leq \alpha^* \rangle}(A)$ with $J_\alpha = id_{\bar{A} \restriction \alpha}$).
4) If $rk'_{\bar{A}}(\kappa) = \zeta$, then we can find $\bar{Y} = \langle Y_\varepsilon : \varepsilon < \zeta \rangle$, an increasing sequence of subsets of α^*, $\bigcup\limits_{\varepsilon < \zeta} Y_\varepsilon = \alpha^*$ and $\{A_\alpha : \alpha \in Y_\varepsilon \backslash \bigcup\limits_{\xi < \varepsilon} Y_\xi\}$ are almost disjoint modulo the ideal generated by $\{A_\alpha : \alpha \in \bigcup\limits_{\xi < \varepsilon} Y_\xi\}$.
5) If $\bar{J} = \langle (J_\alpha, J'_\alpha) : \alpha < \alpha^* \rangle$ is as in 2.1(2), where J_α, J'_α are ideals on $\kappa, J'_\alpha = J_\alpha + A_\alpha$ and $\bar{A} = \langle A_\alpha : \alpha < \alpha^* \rangle$ then for every $B \subseteq \kappa$ we have $rk_{\bar{A}}(B) \geq rk'_{\bar{A}}(B) \geq rk'_{\bar{J}}(B)$.

Proof. Straight: e.g. for the fourth part use $Y_\varepsilon =: \{\alpha : \mathrm{rk}'_{\bar{A}}(A_\alpha) \leq \varepsilon\}$. $\square_{2.2}$

2.3 Claim. *1) If $rk_{\bar{A}}(\kappa) \geq \kappa^+$ then for some $\bar{\alpha} = \langle \alpha_n : n < \omega \rangle$ we have $\alpha_n < \alpha_{n+1} < \kappa$ and for every $\ell < k < \omega$ for some $\alpha < \alpha^*$ we have $A_\alpha \cap \{\alpha_\ell, \alpha_{\ell+1}, \ldots, \alpha_k\} = \{\alpha_{\ell+1}, \ldots, \alpha_k\}$.*
2) If $rk'_{\bar{J}}(\kappa) \geq \beta$ and $\bar{J} = \langle (J_\alpha, J'_\alpha) : \alpha < \alpha^* \rangle$ as in 2.1(2) then for some $\Gamma \subseteq \alpha^*, |\Gamma| \leq |\beta|$ we have $rk'_{\bar{J} \restriction \Gamma}(\kappa) \geq \beta$.
3) If $rk_{\bar{J}}(B) \geq \beta, \bar{J} = \langle (J_\alpha, J'_\alpha) : \alpha < \alpha^* \rangle$ as in 2.1(2) and $J'_\alpha = J_\alpha + A_\alpha$ then we can find $\Gamma \subseteq \alpha^*$ such that

> $(*)$ *$|\Gamma| \leq |\beta| + \aleph_0$ (even $|\Gamma| < |\beta|^+ + \aleph_0$) and if $A_\alpha \subseteq A'_\alpha \in J'_\alpha$ then $rk_{\langle A'_\alpha : \alpha \in J \rangle}(B) \geq \beta$.*

Proof. 1) Let rk $=$ $\mathrm{rk}_{\bar{A}}$; choose by induction on n an ordinal $\alpha_n < \kappa$ and for every $\zeta < \kappa^+$ a decreasing sequence $\langle B^n_{\zeta,0}, \dots, B^n_{\zeta,n} \rangle$ of sets such that

- (α) $(\forall \ell \le n)(\forall m \le n)[\alpha_\ell \in B^n_{\zeta,m} \Leftrightarrow \ell > m]$,
- (β) $\mathrm{rk}(B^n_{\zeta,n}) \ge \zeta$
- (γ) each $B^m_{\zeta,n}$ is the intersection of finitely many A_α's

For $n = 0$, for every $\zeta < \kappa^+$ there is $\alpha_\zeta < \alpha^*$ such that $\kappa \cap A_{\alpha_\zeta} \ne \kappa$, $\mathrm{rk}(\kappa \cap A_{\alpha_\zeta}) \ge \zeta$, and choose $\alpha^0_\zeta \in \kappa \backslash A_{\alpha_\zeta}$. So for some $\alpha_0 < \kappa, \kappa^+ = \sup\{\zeta < \kappa^+ : \alpha^0_\zeta = \alpha_0\}$ and let $B^0_{\zeta,0} = A_{\alpha_{\xi(\zeta)}}$ where $\xi(\zeta) < \kappa^+$ is the minimal $\xi > \zeta$ such that $\alpha^0_\xi = \alpha_0$, as in demand (β) we ask "$\ge \zeta$" not "$= \zeta$", we succeed. If we have defined for n, for each $\zeta < \kappa^+$, as $\mathrm{rk}(B^n_{\zeta+1,n}) \ge \zeta + 1$, there is $\beta(\zeta, n) < \ell g(\bar{A})$ such that $\neg[B^n_{\zeta+1,n} \subseteq A_{\beta(\zeta,n)}]$ but $\mathrm{rk}_{\bar{A}}(B^n_{\zeta+1,n} \cap A_{\beta(\zeta,n)}) \ge \zeta$, and choose $\gamma(\zeta, n) \in B^n_{\zeta+1,n} \backslash A_{\beta(\zeta,n)}$ so for some $\alpha_{n+1} < \kappa$, the set $S_n = \{\zeta < \kappa^+ : \gamma(\zeta, n) = \alpha_{n+1}\}$ is unbounded in κ^+. For every $\zeta < \kappa^+$ let $\xi(\zeta, n) = \min\{\xi : \xi \in S_n, \xi > \zeta\}$, let $B^{n+1}_{\zeta,\ell}$ be $B^n_{\xi(\zeta,n),\ell}$ if $\ell \le n$ and $B^n_{\xi(\zeta,n)+1,n} \cap A_{\beta(\zeta,n)}$ if $\ell = n + 1$. In the end we know that for $\ell < k < \omega$, for every $\zeta < \kappa^+$ we have $B^k_{\zeta,\ell} \cap \{\alpha_\ell, \alpha_{\ell+1}, \dots, \alpha_k\} = \{\alpha_{\ell+1}, \dots, \alpha_k\}$; also $B^k_{\zeta,\ell}$ has the form $\bigcap\limits_{m<m(*)} A_{\alpha(m)}$ for some $\alpha(m) < \ell g(\bar{A})$, so for some m we have $\alpha_\ell \notin A_{\alpha)(m)}$, but $\{\alpha_{\ell+1}, \dots, \alpha_k\} \subseteq B^k_{\zeta,\ell} \subseteq A_{\alpha(m)}$ so $\alpha(m)$ is as required in 2.3(1) for our $\ell < k < \omega$. Lastly $\bigwedge\limits_{n<m} \alpha_n \ne \alpha_m$ hence by Ramsey theorem without loss of generality $\alpha_n < \alpha_{n+1}$ and we are done.
2) By induction on β or by part (3).
3) We can find $\langle (B_\eta, j_\eta) : \eta \in ds(\beta) \rangle$ where $ds(\beta) = \{\eta : \eta$ is a (strictly) decreasing sequence of cardinals $< \beta\}, B_{<>} = B, \mathrm{rk}'_{\bar{J}}(B_\eta) \ge \min(\{\beta\} \cup \{\eta(\ell) : \ell < \ell g(\eta)\}$ and $j_\eta < \alpha^*$ and if $\nu = \eta\hat{\ }\langle\gamma\rangle \in ds(\beta)$ then $B_\nu \ne \beta_\eta, B_\nu = B_\eta \cap A_{j_\eta} \notin J_{j_\eta}$ and $B_\eta \backslash B_\nu \notin J_{j_\eta}$. Let $\Gamma = \{j_\eta : \eta \in ds(\eta)\}$, now if $A_\alpha \subseteq A'_\alpha \in J_\alpha$ for $\alpha \in \Gamma$ then we can prove by induction on $\gamma < \beta$ that: $\eta \in ds(\beta) \Rightarrow \mathrm{rk}'_{\langle A'_\alpha : \alpha \in \Gamma \rangle}(B_\eta) \ge \max(\{\beta\} \cup \{\eta(\ell) : \ell < \ell g(\eta)\}$. $\square_{2.3}$

2.4 Definition. 1) For $\bar{\lambda} = \langle \lambda_n : n < \omega \rangle$ strictly increasing, let $\mathrm{IND}(\bar{\lambda})$ mean:

$(*)_{\bar{\lambda}}$ for every algebra M with universe $\bigcup\limits_{n<\omega} \lambda_n$ and $\aleph_0$ functions (all finitary) there is $\bar{\alpha} = \langle \alpha_n : n < \omega \rangle$ such that:

- (a) $\alpha_n < \lambda_n$
- (b) α_n is not in the M-closure of

$$\{\alpha_\ell : \ell \in (n, \omega)\} \cup \{i : \bigvee_{m<n} i < \lambda_m\}.$$

2) $\mathrm{IND}(\lambda)$ means that $\lambda >$ $\mathrm{cf}(\lambda) = \aleph_0$ and for every (equivalently some, see below) $\bar{\lambda} = \langle \lambda_n : n < \omega \rangle$ strictly increasing with limit λ we have

$(*)'_{\bar{\lambda}}$ for every algebra M with universe λ and countably many functions there is $\langle \alpha_n : n \in \omega \rangle$ such that:

- (a) $w \subseteq w$ is infinite
- (b) $\alpha_n < \lambda_n$ for $n \in w$
- (c) for $n \in w, \alpha_n$ is not in the M-closure of

$$\{\alpha_\ell : \ell \in w, \ell > n\} \cup \{i : \bigvee_{\ell \in n \cap w} i < \lambda_\ell\}$$

3) $\mathrm{IND}(\lambda, \kappa) = \mathrm{IND}^0(\lambda, \kappa)$ means: if M is a model with universe λ and κ functions we can find $\bar{\alpha} = \langle \alpha_n : n < \omega \rangle$ such that

$$\alpha_n < \lambda, \alpha_n \notin c\ell_M\{\alpha_\ell : \ell < \omega, \ell > n\}.$$

4) $\mathrm{IND}^1(\lambda, \kappa)$ is defined similarly but demanding

$$\alpha_n \notin c\ell_M\{\alpha_\ell : e, s, p, r, < \omega, \ell \neq n\}.$$

2.5 Observation. 1) In 2.4(2), if $(*)'_{\bar{\lambda}}$ holds for one $\bar{\lambda}$ with limit λ, then it holds for every $\bar{\lambda}' = \langle \lambda'_n : n < \omega \rangle$ with limit λ.
2) If λ is uncountable with cofinality $\aleph_0, P$ a forcing notion of cardinality $\leq \mu < \lambda$ or satisfying the μ^+-c.c. for some $\mu < \lambda$, or λ-complete then: $\mathrm{IND}(\lambda) \Leftrightarrow \Vdash_P$ "$\mathrm{IND}(\lambda)$" and if $\kappa \in [\mu, \lambda), \mu$ as above then $\mathrm{IND}(\lambda, \kappa) \Leftrightarrow \Vdash_P$ "$\mathrm{IND}(\lambda, \kappa)$" and if in addition $\mu < \lambda_n < \lambda_{n+1}$, then $\mathrm{IND}(\langle \lambda_n : n < \omega \rangle) \Leftrightarrow \Vdash_P$ " $\mathrm{IND}(\langle \lambda_n : n < \omega \rangle)$".
3) $\mathrm{IND}(\bar{\lambda}) \Rightarrow \mathrm{IND}(\sum_{n<\omega} \lambda_n) \Rightarrow \mathrm{IND}^1(\lambda, \kappa) \Rightarrow \mathrm{IND}^0(\lambda, \kappa)$ if $\lambda = \bigcup_{n<\omega} \lambda_n$ and $\lambda_n < \lambda_{n+1}$ and $\lambda_0 > \kappa$. If $\kappa < \lambda \leq \lambda'$ then

$$\mathrm{IND}^i(\lambda, \kappa) \Rightarrow \mathrm{IND}^i(\lambda', \kappa).$$

4) If ($i \in \{0, 1\}$ and) $\mathrm{IND}^i(\lambda, \kappa), \lambda$ minimal for this κ then

- (a) $\kappa \leq \kappa_1 < \lambda \Rightarrow \mathrm{IND}^i(\lambda, \kappa_1)$
- (b) $\mathrm{cf}(\lambda) = \aleph_0$ and $\mathrm{IND}^1(\lambda, \kappa)$ or λ is inaccessible
- (c) if $\lambda = \sum_{n<\omega} \lambda_n$ and $\lambda_n < \lambda_{n+1}$ then not only $(*)'_{\bar{\lambda}}$ (where $\bar{\lambda} = \langle \lambda_n : n < \omega \rangle$) but if P is a c.c.c. forcing adding a dominating real then in V^P for some infinite $w \subseteq \omega$ we have $(*)_{\bar{\lambda} \restriction w}$ holds.

5) $\mathrm{IND}^1(\lambda, \kappa)$ is equivalent to $\mathrm{IND}^0(\lambda, \kappa)$.

Proof. 1, 2), 3) Check.
4) Clause (a) of (a): Assume not and first let $i = 0$. Let $\chi = \beth_3(\lambda)^+$ and let M be the model with universe λ and the functions (n-place from λ to λ for some n) definable in $(\mathscr{H}(\chi), \in, <^*_\chi)$ with the parameters $\lambda, \kappa, \kappa_1$. Clearly $(M, \alpha)_{\alpha < \kappa_1}$

exemplifies $\neg\mathrm{IND}^i(\lambda,\kappa_1)$. Let F^-_n, F^+_n be such that for $\bar\beta = \langle \beta_\ell : \ell < n\rangle, \beta_\ell < \lambda$ we have: $F^+_n(-,\bar\beta)$ is a one-to-one function from $c\ell_M(\{\beta_0,\dots,\beta_{n-1}\}\cup\kappa_1)$ onto κ_1 and $F^-_n(-,\beta)$ is its inverse. We can apply the assumption $\mathrm{IND}^i(\lambda,\kappa)$ to the model $(M, F^+_n, F^-_n, \beta)_{n<\omega,\beta<\kappa}$, so there are $\alpha_n (n<\omega)$ as in 2.4(3). By the assumption for no infinite $w \subseteq \omega$ is $\{\alpha_n : n \in w\}$ as required in 2.4(3) for $(M,\beta)_{\beta<\kappa_1}$ hence for some infinite $w\subseteq\omega$, $\bigwedge\limits_{n\in w} \alpha_n \in c\ell_M(\{\alpha_\ell : n < \ell \in w\}\cup\kappa_1)$, just choose by induction on $\ell<\omega, u_\ell, n_\ell$ such that: $\bigwedge\limits_{m<\ell} n_m < n_\ell < \omega, u_\ell \subseteq (n_\ell,\omega)$ is infinite, $u_{\ell+1}\subseteq u_\ell\subseteq w$ and $\alpha_{n_\ell}\notin c\ell_M(\{\alpha_n : n\in u_\ell\})$, we cannot succeed so w or some u_ℓ is as required. By renaming $w=\omega$, so for every n for some $k_n \in (n+1,\omega)$ we have $\alpha_n \in c\ell)M(\{\alpha_{n+1},\dots,\alpha_{k_n}\}\cup\kappa_1)$; as we can increase k_n, without loss of generality $k_n < k_{n+1}$ hence $m\le n \Rightarrow \alpha_m \in c\ell_M(\{\alpha_{n+1},\dots,\alpha_{k_n}\}\cup\kappa_1)$ (just prove this by induction on n), so $\gamma_{n,m} =: F^+_{k_n-n}(\alpha_m,\alpha_{n+1},\dots,\alpha_{k_n}) < \kappa_1$ and for each n we have: $\langle\gamma_{n,m} : m\le n\rangle$ is with no repetitions. Choose by induction on $\ell, m_\ell \in [\ell^2,(\ell+1)^2)$ such that $\gamma_{(\ell+1)^2,m_\ell}\notin\{\gamma_{(q+1)^2,m_q} : q<\ell\}$. But as $\neg$ $\mathrm{IND}^i(\kappa_1,\kappa)$ (because $\lambda>\kappa_1$ was minimal such that ...) for some $\ell<p<\omega$ we have $\gamma_{(\ell+1)^2,m_\ell}\in c\ell_M(\{\gamma_{(q+1)^2,m_q} : \ell<q<p\}\cup\kappa)$ and using some $F^-_{p^2-(\ell+1)^2}$ we have $\alpha_{m_\ell}\in c\ell_M(\{\alpha_q : q$ is $\ge (\ell+1)^2$ but $\le k_{(p^2)}\}\cup\kappa)$.
[Why? First note that $\gamma_{(q+1)^2,m_q}$ belong to this model for $q=\ell+1,\dots,p-1$, (using $F_{k_{(q+1)^2}}$) hence also $\gamma_{(\ell+1)^2,m_\ell}$ belongs to this model by the choice of ℓ and p; a contradiction.]

If $i=1$ the proof is similar: choose, by induction on $\ell, k_\ell, m_\ell, m_\ell$ such that $k_\ell<m_\ell<k_{\ell+1}$ and $\alpha_{m_\ell}\in c\ell_M(\{\alpha_n : n\in[k_\ell,m_\ell)$ or $n\in(m_\ell,k_{\ell+1})\}\cup\kappa_1)$, this is possible as otherwise $\{\alpha_n : n\in[k_\ell,\omega)\}$ contradict "M exemplifies $\neg\mathrm{IND}^1(\lambda,\kappa_1)$". Let

$$\gamma_\ell = F^+_{k_{\ell+1}-k_\ell-1}(\alpha_{m_\ell};\alpha_{k_\ell},\alpha_{k_\ell+1},\dots,\alpha_{m_\ell-1},\alpha_{m_\ell+1},\dots,\alpha_{k_{\ell+1}-1}) < \kappa_1.$$

For some $\ell<\ell(*)<\omega$ we have $\gamma_\ell\in c\ell_M(\{\gamma_0,\dots,\gamma_{\ell-1},\gamma_{\ell+1},\dots,\gamma_{\ell(*)-1}\}\cup\kappa)$ (because $\neg$ $\mathrm{IND}^1(\kappa_1,\kappa)$ as λ is first and the choice of M), hence $\alpha_\ell\in c\ell_M(\{\alpha_n : n<\omega, n\le\ell\}\cup\kappa)$; a contradiction.

Clause (b) of (4): By the definition easily $\aleph_0 < \mathrm{cf}(\lambda)\le\kappa$ is impossible.
[Why? Let $\lambda=\sum\limits_{i<\kappa}\lambda_i$, with $\lambda_i<\lambda$, and by the minimality of λ let M_i be a model with universe λ_i and $\le\kappa$ functions exemplifying $\neg\mathrm{IND}(\lambda_i,\kappa)$ and lastly let M be the model with universe λ and the functions of all the M_i; check that M exemplifies $\neg$ $\mathrm{IND}(\lambda,\kappa)$.]
By 2.5(4) clause (a) it follows that

$$[\mathrm{cf}(\lambda)>\aleph_0 \ \& \ \kappa_1<\lambda\Rightarrow\kappa_1<\mathrm{cf}(\lambda)].$$

So if $\mathrm{cf}(\lambda)>\aleph_0$ then λ is regular, it is inaccessible as it is not a successor as trivially

$$\neg\mathrm{IND}^i(\mu,\kappa)\Rightarrow\neg\mathrm{IND}(\mu^+,\kappa).$$

We still have to prove $\mathrm{IND}^1(\lambda)$ when $\mathrm{cf}(\lambda) = \aleph_0$; if $i = 1$ this is trivial so assume $i = 0$. So assume $\mathrm{cf}(\lambda) = \aleph_0, \bar\lambda = \langle \lambda_n : n < \omega\rangle, \lambda_n < \lambda_{n+1}$ and $\lambda = \sum\limits_{n<\omega} \lambda_n$. We should prove $\mathrm{IND}^1(\lambda,\kappa)$, so let us be given M as in 2.4(4). Without loss of generality, expanding M by examples to $\neg\mathrm{IND}(\lambda_n,\kappa)$, we get M^+.
As $\mathrm{IND}^0(\lambda,\kappa)$ there are $\alpha_n < \lambda$ (for $n < \omega$) such that $\alpha_n \notin c\ell_{M^+}\{\alpha_\ell : \ell < \omega, \ell > n\}$. Without loss of generality $\langle \alpha_\ell : \ell < \omega\rangle$ is strictly increasing, and letting $m_n = \min\{m : \lambda_m > \alpha_n\}$, without loss of generality $\langle m_n : n < \omega\rangle$ is constant or strictly increasing. If $\bigwedge\limits_n M - n = m(*)$ we get a contradiction to "M^+ expand a counterexample to $\mathrm{IND}^0(\lambda_{m(*)},\kappa)$", so $\bigwedge\limits_n m_n < m_{n+1}$. It suffices to prove that for every $n_1 < \omega$ there is $n_2 \in (n_1,\omega)$ such that $\alpha_{n_2} \notin c\ell_M(\{\alpha_\ell : \ell < \omega, \ell > n_2\} \cup \lambda_{n_1})$ as then we can produce an example for $(*)'_{\bar\lambda}$. But if for some n_1 there is no n_2 as above the proof of clause (a) gives a contradiction.

Clause (c) of (4): Left to the reader.
5) By 2.5(3),

$$\mathrm{IND}^1(\lambda,\kappa) \Rightarrow \mathrm{IND}^0(\lambda,\kappa) \text{ and } \lambda \le \lambda' \ \ \& \ \ \mathrm{IND}^i(\lambda,\kappa) \Rightarrow \ \mathrm{IND}^i(\lambda,\kappa).$$

Hence it suffices to prov: if λ is minimal such that $\mathrm{IND}^0(\lambda,\kappa)$ then $\mathrm{IND}^1(\lambda,\kappa)$. let M be a model with universe λ and vocabulary of cardinality $\le \kappa$ and we shall prove the conclusion of 2.4(4) (= the Definition of $\mathrm{IND}^1(\lambda,\kappa)$). Let for $n < \omega, F^+_n, F^-_n$ be $(n+2)$-place functions from λ to λ such that

- $(*)$ if $\gamma < \lambda$ and $\bar\beta \in {}^n\lambda$ then $F^+_n(-,\bar\beta,\gamma)$ is a one-to-one function from $c\ell_M(\{\beta_\ell : \ell < \ell g(\bar\beta)\} \cup \{i : i \le \gamma \text{ or } i < \kappa\})$ onto $|\kappa+\gamma|$ and $F^-_n(-,\bar\beta,\gamma)$ be the inverse function.

Let $M^* = (M, F^+_n, F^-_n)_{n<\omega}, M^+ = (M^*, i)_{i<\kappa}$ but as $\mathrm{IND}(\lambda,\kappa)$ we can apply Definition 2.4(4) and get $\langle \alpha_n : n < \omega\rangle, \alpha_n < \lambda, \alpha_n \notin c\ell_{M^+}(\{\alpha_\ell : \ell \in (n,\omega)\})$. Without loss of generality $\langle \alpha_n : n < \omega\rangle$ is strictly increasing; $\alpha_n > \kappa$ (as without loss of generality each $i \le \kappa$ is an individual constant of M), clearly it suffices to prove

- $(**)$ for any $n < \omega$ for some $m \in (n,\omega)$,
  $\alpha_m \notin c\ell_M(\{\alpha_\ell : \ell < n \text{ or } \ell > m\})$ hence it suffices to prove
- $(**)'$ for any $n < \omega$ for some $m \in (n,\omega)$,

$$\alpha_m \notin c\ell_M(\{\alpha_\ell : \ell > m\} \cup \{i : i \le \alpha_m\}).$$

But here we can repeat the proof of clause (a) of 2.4(4) for the case $i = 0$. $\square_{2.5}$

2.6 Claim. *1) Assume* $\lambda >\ cf(\lambda), |\mathfrak{a}| < \min(\mathfrak{a})$ *and* $sup(\mathfrak{a}) = \lambda$ *and* $rk'_{\langle J_{<\theta}[\mathfrak{a}]:\theta\in\mathrm{pcf}(\mathfrak{a})\rangle}(\mathfrak{a}) \ge |\mathfrak{a}|^+$. *Then* $IND(\lambda,|\mathfrak{a}|)$.
2) Moreover, for any model M *with universe* λ *and* $|\mathfrak{a}|$ *functions and* $\mathfrak{c}$ *such that* $\mathfrak{a} \subseteq \mathfrak{c} \subseteq\ pcf(\mathfrak{a}), |\mathfrak{c}| < \min(\mathfrak{a})$ *and* $\langle \mathfrak{b}_\theta[\mathfrak{a}] : \theta \in\ pcf(\mathfrak{a})\rangle$ *a generating sequence we can find* $\bar\alpha = \langle \alpha_\theta : \theta\mathfrak{a}\rangle \in \Pi\mathfrak{a}$ *such that, defining for* $\mathfrak{b} \subseteq \mathfrak{a}$*:*

$$c\ell_{M,\bar\alpha}(\mathfrak{b}) = \mathfrak{b} \cup \{\theta \in \mathfrak{a} : \alpha_\lambda \in c\ell_M(\{\alpha_\mu : \mu \in \mathfrak{b}\})\};$$

we have

$\otimes_1$ $[\theta \in \mathfrak{c} \Rightarrow c\ell_{M,\bar\alpha}(\mathfrak{b}_\theta[\mathfrak{a}]) \in J_{\leq\theta}[\mathfrak{a}]]$*;*

$\otimes_2$ $c\ell_{M,\bar\alpha}(-)$ *is a closure operation on* $\mathfrak{a}$*, i.e.*

$$\mathfrak{b}_1 \subseteq \mathfrak{b}_2 \Rightarrow c\ell_{M,\bar\alpha}(\mathfrak{b}_1) \subseteq c\ell_{M,\bar\alpha}(\mathfrak{b}_2),$$

$$\mathfrak{b} \subseteq c\ell_{M,\alpha}(\mathfrak{b}),$$

$$c\ell_{M,\bar\alpha}(c\ell_{M,\bar\alpha}(\mathfrak{b})) = c\ell_{M,\bar\alpha}(\mathfrak{b}).$$

Remark. See 3.14 - 3.17 for more.

Proof. 1) Let us define $\bar J = \langle (J_{<\theta}[\mathfrak{a}], J_{\leq\theta}[\mathfrak{a}]) : \theta \in \text{pcf}(\mathfrak{a})\rangle$. We prove part (1) assuming part (2). Choose $\mathfrak{c} \subseteq \text{pcf}(\mathfrak{a}), |\mathfrak{c}| = |\mathfrak{a}|^+$ such that $\text{rk}'_{\bar J \restriction \mathfrak{c}}(\mathfrak{a}) \geq |\mathfrak{a}|^+$ (this is possible by 2.3(2) and without loss of generality $\mathfrak{a} \subseteq \mathfrak{c}$ and let $\langle \mathfrak{b}_\theta[\mathfrak{a}] : \theta \in \text{pcf}(\mathfrak{a})\rangle$ be a generating sequence for $\mathfrak{a}$ (exists by [Sh 371, 2.6]). For proving IND$(\lambda, |\mathfrak{a}|)$ let M be a model with universe λ and $\leq |\mathfrak{a}|$ functions, by part (2) there is a sequence $\bar\lambda = \langle \alpha_\tau : \tau \in \mathfrak{c}\rangle$ as there. Let for $\theta \in \mathfrak{c}, \mathfrak{d}_\theta =: c\ell_{M,\bar\alpha}(\mathfrak{b}_\theta[\mathfrak{a}]) \subseteq \mathfrak{a}$ (as defined in part (2)) so $\mathfrak{b}_\theta[\mathfrak{a}] \subseteq \mathfrak{d}_\theta \in J_{\leq\theta}[\mathfrak{a}]$ hence $J_{<\theta}[\mathfrak{a}] + \mathfrak{d}_\theta = J_{\leq\theta}[\mathfrak{a}]$. So by 2.2(5) we know $\text{rk}_{\langle \mathfrak{d}_\theta : \theta \in \mathfrak{c}\rangle}(\mathfrak{a}) \geq \text{rk}'_{\langle \mathfrak{d}_\theta : \theta \in \mathfrak{c}\rangle}(\mathfrak{a}) \geq \text{rk}'_{\bar J \restriction \mathfrak{c}}(\mathfrak{a}) \geq \kappa^+$ (by the choice of $\mathfrak{c}$ above). Now by 2.3(1) we can find $\tau_n \in \mathfrak{a}$ for $n < \omega$, pairwise distinct and strictly increasing with n such that for every $n < m < \omega$ for soe $\theta_{n,m} \in \mathfrak{c}$ we have $\{\tau_n, \tau_{n+1}, \dots, \tau_m\} \cap \mathfrak{d}_{\theta_{n,m}} = \{\tau_{n+1}, \dots, \tau_m\}$, note: as $\tau_m \in \mathfrak{d}_{\theta_{n,m}}$ necessarily $\theta_{n,m} \geq \tau_m$. So by the choice of $\bar\alpha$, we have $\alpha_{\tau_n} \notin c\ell_M(\{\alpha_{\tau_{n+1}}, \alpha_{\tau_{n+2}}, \dots, \alpha_{\tau_m}\})$. So $\langle \alpha_{\tau_n} : n < \omega\rangle$ are as required in the definition of IND$(\lambda, |\mathfrak{a}|)$.

2) Let $\bar{\mathfrak{b}} = \langle \mathfrak{b}_\theta[\mathfrak{a}] : \theta \in \text{pcf}(\mathfrak{a})\rangle$ be the generating sequence for $\mathfrak{a}$; without loss of generality max pcf$(\mathfrak{a}) \in \mathfrak{c}$ and $\theta \in \mathfrak{c} \backslash \{\max \text{pcf}(\mathfrak{a})\} \Rightarrow \min(\text{pcf}(\mathfrak{a}) \backslash \theta^+) \in \mathfrak{c}$. We know that

$(*)_{\mathfrak{a}}$ there is $\bar F = \langle F_\theta : \theta \in \text{pcf}(\mathfrak{a})\rangle$ such that:

- (a) $F_\theta \subseteq \Pi\mathfrak{a}$ and $|F_\theta| \leq \theta$
- (b) $\{f \restriction \mathfrak{b} : f \in F_\theta\}$ is cofinal in $\Pi\mathfrak{b}$ for every $\mathfrak{b} \in J_{\leq\theta}[\mathfrak{a}]$
- (c) F_θ includes $\cup\{F_\tau : \tau \in \theta \cap \text{pcf}(\mathfrak{a})\}$ and is closed under some natural operations
- (d) if $\tau \in \theta \cap \mathfrak{c}$ then $f \in F_\theta \Rightarrow (\exists g \in F_\tau)(f \restriction \mathfrak{b}_\tau[\mathfrak{a}] = g \restriction \mathfrak{b}_\tau[\mathfrak{a}])$.

[Why? E.g. the proof of [Sh 355],3.5.]

Let M be a model with universe λ and vocabulary of cardinality $|\mathfrak{a}|$. For every $f \in \Pi\mathfrak{a}$ (e.g., $f \in F_\theta, \theta \in \text{pcf}(\mathfrak{a})$) we define $g_f \in \Pi\mathfrak{a}$ by $g_f(\tau) =: \sup[\tau \cap c\ell_M(\text{Rang } f)]$. For every $\theta \in \mathfrak{c} \cap \text{pcf}(\mathfrak{a}), \{g_f : f \in \bigcup\limits_{\tau \in \theta \cap \mathfrak{c}} F_\tau\}$ is a subset of $\Pi\mathfrak{a}$ of cardinality $< \theta$ (here instead $|\mathfrak{c}| < \min(\mathfrak{a})$, just $\theta \in \mathfrak{c} \Rightarrow |\mathfrak{c} \cap \theta| < \theta$ suffice), so there is $g^\theta \in F_\theta$ such that:

$(*)_1$ $f \in \bigcup\limits_{\tau \in \theta \cap \mathfrak{c}} F_\tau \Rightarrow g_f < g^\theta \bmod J_{<\theta}[\mathfrak{a}]$.

Define $g^* \in \Pi\mathfrak{a}$ by $g^*(\tau) = \sup\{g^\theta(\tau) + 1 : \theta \in \mathfrak{c}\}$ (remember $|\mathfrak{c}| < \min(\mathfrak{a})$). So there is $h \in F_{\max \text{ pcf}(\mathfrak{a})}$ such that $g^* < h$ (see $(*)_{\mathfrak{a}}(b)$); we shall show that:

$(*)_2$ for any such h the sequence $\langle h(\tau) : \tau \in \mathfrak{a}\rangle$ is as required.

Proof of $(*)_2$. So let $\alpha_\tau = h(\tau)$. Note that $\otimes_2$ from 2.6(2) is trivial so we shall prove $\otimes_1$. Assume $\theta \in \mathfrak{c}$ and let $\mathfrak{b} = \mathfrak{b}_\theta[\mathfrak{a}] \in J_{\leq\theta}[\mathfrak{a}]$ so by clause (d) of $(*)_{\mathfrak{a}}$ for some $f_1 \in F_\theta$ we have

$\oplus_1$ $h \restriction \mathfrak{b} = f_1 \restriction \mathfrak{b}$;

we can assume $\theta <\ \max \text{pcf}(\mathfrak{a})$ (otherwise conclusion is trivial) and let $\sigma = \min(\text{pcf}(\mathfrak{a}) \backslash \theta^+)$, by an assumption made in the beginning of the proof $\sigma \in \mathfrak{c}$ and so as $f_1 \in F_\theta$ by the choice of g^σ we have:

$$g_{f_1} < g^\sigma \bmod J_{<\sigma}[\mathfrak{a}]$$

but by the choice of g^*

$$g^\sigma < g^*$$

and by the demand of h

$$g^* < h$$

together

$$g_{f_1} < h \bmod J_{<\sigma}[\mathfrak{a}]$$

so for some $\mathfrak{d} \in J_{<\sigma}[\mathfrak{a}] = J_{\leq\theta}[\mathfrak{a}]$ we have:

$\oplus_2$ $g_{f_1} \restriction (\mathfrak{a} \backslash \mathfrak{d}) < h \restriction (\mathfrak{a} \backslash \mathfrak{d})$.

Now for any $\tau \in \mathfrak{b}$

$\oplus_3$ $\tau \in c\ell_{M,\bar{\alpha}}(\mathfrak{b}) \Rightarrow \alpha_\tau \in c\ell_M[\{\alpha_\kappa : \kappa \in \mathfrak{b}\}] \Rightarrow \alpha_\tau \in c\ell_M[\text{Rang}(h \restriction \mathfrak{b})] \Rightarrow \alpha_\tau \in c\ell_M[(\text{Rang}(f_1 \restriction \mathfrak{b})] \Rightarrow \alpha_\tau \in c\ell_M(\text{Rang}(f_1)) \Rightarrow \alpha_\tau \leq g_{f_1}(\tau)$.

By $\oplus_2 + \oplus_3$ the required conclusion follows. $\square_{2.6}$

E.g.

2.7 Claim. *If* $pp(\aleph_\omega) > \aleph_{\omega_1}$ *then* $IND(\aleph_\omega)$.

Proof. Let $\mathrm{pp}(\aleph_\omega) = \aleph_{\alpha^*}$ (so $\alpha^* < \omega_4, \alpha^* = \beta^* + 1$, see [Sh:g, Ch.IX,2.1]) let $\mathfrak{a} = \{\aleph_{i+1} : 5 \le i < \omega\}$ so we know $\mathrm{pcf}(\mathfrak{a}) = \{\aleph_{i+1} : 5 \le i < \alpha^*\}$ and let $\bar{\mathfrak{b}} = \langle \mathfrak{b}_\lambda : \lambda \in \mathrm{pcf}(\mathfrak{a})\rangle$ be a normal generating sequence for $\mathrm{pcf}(\mathfrak{a})$ (not $\mathfrak{a}$!); without loss of generality $\mathfrak{b}_{\aleph_{\alpha^*}} = \mathrm{pcf}(\mathfrak{a})$. Let $\bar{\mathfrak{c}} = \langle \mathfrak{b}_\lambda \cap \mathfrak{a} : \lambda \in \mathrm{pcf}(\mathfrak{a})\rangle$. Now by localization ([Sh:g, Ch.VIII], 2.6) we know that for some club E of $\omega_1 : \delta \in E \Rightarrow \mathrm{pp}(\aleph_\delta) > \aleph_{\omega_1}$ (so we can assume $\omega \in E$).

Let $E = \{\beta_\zeta : \zeta < \omega_1\}$ (increasing in ζ). Hence by [Sh:g, Ch.II], 1.5A we have

$(*)_0$ for limit $\delta < \omega_1, \delta < \beta < \omega_1, \delta \in E$, for some unbounded $\mathfrak{d} \in \aleph_\delta \cap$ Reg, we have $\aleph_{\beta+1} \in \mathrm{pcf}_{J^{\mathrm{bd}}_{\mathfrak{d}}}(\mathfrak{d})$.

Hence we can prove by induction on $\varepsilon < \omega_1$ that:

$(*)$ if $\beta_\varepsilon \le \zeta < \omega_1, \mathfrak{b} \subseteq \mathfrak{a}$ and $\mathfrak{b} = \mathfrak{a} \cap \mathfrak{b}_{\aleph_{\zeta+1}} \bmod J_{<\aleph_{\zeta+1}}[\mathfrak{a}]$ then $\mathrm{rk}'_{\bar{\mathfrak{c}}}(\mathfrak{b}) \ge \varepsilon$.

[Why? For $\varepsilon = 0$ this is trivial and also for ε limit. If $\varepsilon = \xi + 1$ we can find $\mathfrak{d} \in \aleph_{\beta_\varepsilon} \cap \mathrm{Reg} \setminus \aleph_{\beta_\xi}$ such that $\aleph_{\zeta+1} \in \mathrm{pcf}_{J^{\mathrm{bd}}_{\mathfrak{d}}}$ so without loss of generality $\aleph_{\zeta+1} = \max \mathrm{pcf}(\mathfrak{d})$. By [Sh:g, Ch.I], **?** we have that the set $\mathfrak{d}'$ of $\theta \in \mathfrak{d}$ such that $\mathfrak{b}_\theta \cap$
—> scite{1.12} undefined
$\mathfrak{a} \subseteq \mathfrak{b} \bmod J_{<\theta}[\mathfrak{a}]$ is $= \mathfrak{d} \bmod J_{<\aleph_{\zeta+1}}[\mathfrak{d}]$ hence is not bounded in $\mathfrak{d}$, hence is not bounded in $\aleph_{\beta_\varepsilon}$. But $\mathfrak{d}' \subseteq \mathfrak{d} \subseteq \aleph_{\beta_\varepsilon} \cap \mathrm{Reg} \setminus \aleph_{\beta_\varepsilon}$, hence by the induction hypothesis $\theta \in \mathfrak{d}' \Rightarrow \mathrm{rk}'_{\bar{\mathfrak{c}}}(\mathfrak{b}_\theta \cap \mathfrak{b}) \ge \xi$, but of course $\mathfrak{b} \setminus \mathfrak{b}_\theta \ne \emptyset$.]

Now apply 2.6. $\square_{2.7}$

2.8 Proclaim. *If* $|\mathfrak{a}| < min(\mathfrak{a}), \lambda = \sup(\mathfrak{a})$ *is singular and* $pcf_{J^{\mathrm{bd}}_{\mathfrak{a}}}(\mathfrak{a})$ *contains an interval of* Reg *of cardinality* $|\mathfrak{a}|^+$ *then* $IND(\lambda)$.

Proof. Similar to the proof of 2.7.

2.9 Discussion. We can also prove e.g.: if $\lambda = \mathrm{tcf}(\prod_{\varepsilon<\kappa} \lambda_\varepsilon / [\kappa]^{<\aleph_0})$, satisfies $\lambda > \lambda_\varepsilon = \mathrm{cf}(\lambda_\varepsilon) > \kappa$, then for every algebra M on $\sum_{\varepsilon<\kappa} \lambda_\varepsilon$ with $< \min\{\lambda_\varepsilon : \varepsilon < \kappa\}$ functions there are $\alpha_\varepsilon < \lambda_\varepsilon (\varepsilon < \kappa)$ such that: for finite $u \subseteq \kappa$ we have $\{\zeta : \alpha_\zeta \in c\ell_M(\{\alpha_\varepsilon : \varepsilon \in u\})\}$ is finite (and more). Not clear how interesting is this statement and where it leads.

2.10 Claim. *Assume* $\mathfrak{a} \subseteq \mathfrak{c} \subseteq pcf(\mathfrak{a})$ *and* $|\mathfrak{c}| < \min(\mathfrak{a})$ *(or* $\mathfrak{c} \in J_*[pcf(\mathfrak{a})]$*, see [Sh:g, Ch.VIII,§3] and [Sh:E11]), so* $pcf(\mathfrak{a}) = pcf(\mathfrak{c})$*. Then*

$$rk'_{\langle J_{<\theta}[\mathfrak{a}]:\theta\in\mathrm{pcf}(\mathfrak{a})\rangle}(\mathfrak{a}) = rk'_{\langle J_{<\theta}[\mathfrak{c}]:\theta\in\mathrm{pcf}(\mathfrak{a}')\rangle}(\mathfrak{c}).$$

Proof. Let $\bar{\mathfrak{b}} = \langle \mathfrak{b}_\theta[\mathfrak{a}] : \theta \in \mathrm{pcf}(\mathfrak{a})\rangle$ be a generating sequence for $\mathfrak{a}$, hence we know that letting $\mathfrak{b}_\theta[\mathfrak{c}] =: \mathfrak{c} \cap \mathrm{pcf}(\mathfrak{b}_\theta[\mathfrak{a}])$, we have: $\bar{\mathfrak{b}}' = \langle \mathfrak{b}_\theta[\mathfrak{c}] : \theta \in \mathrm{pcf}(\mathfrak{a})\rangle$ is a generating sequence for $\mathfrak{c}$. Without loss of generality $\mathfrak{b}_\theta[\mathfrak{c}] \cap \mathfrak{a} = \mathfrak{b}_\theta[\mathfrak{a}]$ and $\mathfrak{b}_{\max\ \mathrm{pcf}(\mathfrak{a})}[\mathfrak{a}] = \mathfrak{a}$ hence $\mathfrak{b}_{\max\ \mathrm{pcf}(\mathfrak{a})}[\mathfrak{c}] = \mathfrak{a}$. So we can prove easily:

$$\mathfrak{d} \subseteq \mathfrak{a} \Rightarrow \ \mathrm{rk}'_{\langle J_{<\theta}[\mathfrak{a}]:\theta \in \mathrm{pcf}(\mathfrak{a})\rangle}(\mathfrak{d}) \leq \ \mathrm{rk}'_{\langle J_{<\theta}[\mathfrak{c}]:\theta \in \mathrm{pcf}(\mathfrak{a})\rangle}(\mathfrak{d})$$

(as $J_{<\theta}[\mathfrak{a}] = J_{<\theta}[\mathfrak{c}] \restriction \mathfrak{a}$).

For the other direction we prove

$(*)$ if $n < \omega, \{\theta_1, \dots, \theta_n\} \subseteq \ \mathrm{pcf}(\mathfrak{a})$ then

$$\mathrm{rk}'_{\langle J_{<\theta}[\mathfrak{c}]:\theta \in \mathrm{pcf}(\mathfrak{a})\rangle}(\bigcap_{\ell=1}^{n} \mathfrak{b}_{\theta_\ell}[\mathfrak{c}]) \leq \ \mathrm{rk}'_{\langle J_{<\theta}[\mathfrak{c}]:\theta \in \mathrm{pcf}(\mathfrak{a})\rangle}(\bigcap_{\ell=1}^{n} \mathfrak{b}_{\theta_\ell}[\mathfrak{a}]).$$

$\square_{2.10}$

§3 EXISTENCE OF FREE SETS IMPLIES RESTRICTIONS ON PCF

3.1 Definition. Suppose $\bar{\mathscr{I}} = \langle (\kappa_n, I_n) : n < \omega \rangle$ is such that: I_n is an ideal on κ_n.
1) We define $J_n = J_n^{\bar{\mathscr{I}}}$ an ideal on $\prod\limits_{\ell < n} \kappa_\ell$:

$(*)_0$ $J_0 =$ the empty ideal on $\{<>\}$

$(*)_1$ $J_{n+1} = \{A \subseteq \prod\limits_{\ell \le n} \kappa_\ell : \{\alpha < \kappa_n : \{\eta \in \prod\limits_{\ell < n} \kappa_\ell : \eta\hat{\ }\langle \alpha \rangle \in A\} \notin J_n\} \in I_n\}$

we let $\bar{J}^{\bar{\mathscr{I}}} = \langle J_n^{\bar{\mathscr{I}}} : n < \omega \rangle$.
2) We say $\langle J_n : n < \omega \rangle$ is a candidate (for $\bar{\mathscr{I}}$) if in $(*)_1$ we weaken "$J_{n+1} = \dots$" to "$J_n \subseteq \dots$". [So there may be many candidates.]

3.2 Fact. 1) In definition 3.1(1) above, each J_n is an ideal on $\prod\limits_{\ell < n} \kappa_\ell$.
2) If $I_0, \dots, I_{n-1}$ are σ-complete then so is J_n.
3) If $\langle J'_n : n < \omega \rangle$ is a candidate for $\bar{\mathscr{I}}$, then $\bigwedge\limits_{n<\omega} J'_n \subseteq J_n^{\bar{\mathscr{I}}}$.

3.3 Claim. *For $\bar{\mathscr{I}}, \bar{J}^{\bar{\mathscr{I}}}$ as in 3.1(1), $A \in J_n$ iff for some functions $f_0, \dots, f_{n-1}$ we have:*

$$Dom(f_\ell) = \prod_{m=\ell+1}^{n-1} \kappa_m \text{ and } Rang(f_\ell) \subseteq I_\ell,$$

and $A \subseteq \bigcup\limits_{\ell < n} A_\ell^n(f_\ell)$ where

$$A_\ell^n(f_\ell) = \{\eta \in \prod_{m<n} \kappa_m : \eta(m) \in f_\ell(\eta \restriction (m, n))\}.$$

Proof. By induction on n.

3.4 Theorem. *Let $\bar{\mathscr{I}} = \langle (\kappa_n, I_n) : n < \omega \rangle$ be as in 3.1, $\kappa = \sum\limits_{n<\omega} \kappa_n \le \mu_0 = cf(\mu_0) < \mu_1 < \mu = cf(\mu)$. Then $\otimes_1 \Rightarrow \otimes_2$ where*

$\otimes_1$ *for each n there is $\langle \lambda_i^n : i < \kappa_n \rangle$ such that:*
$\lambda_i^n \in [\mu_0, \mu_1) \cap$ Reg and $\prod\limits_{i < \kappa_n} \lambda_i^n / I_n$ is μ-directed

$\otimes_2$ *for some $\bar{\mathscr{I}}$-candidate, $\bar{J} = \langle J_n : n < \omega \rangle$ (except for clause (δ), $J_n = J_n^{\bar{\mathscr{I}}}$ is O.K.) we have:*

$\otimes^2_{\bar{J}}$ *there are* $\bar{\lambda}^n = \langle \lambda_\eta : \eta \in \prod\limits_{\ell<n} \kappa_\ell \rangle$ *for* $n \in (0,\omega)$ *such that for each* n*:*

- (α) $\Big(\prod\limits_{\eta \in \prod\limits_{\ell<n} \kappa_\ell} \lambda_\eta\Big)/J^{\bar{\mathscr{I}}}$ *has true cofinality* μ
- (β) $\mu_0 \le \lambda_\eta = \ cf(\lambda_\eta) < \mu_1$ *(note that by clause* (α) *we have* $\{\eta \in \prod\limits_{\ell<n} \kappa_\ell : \lambda_\eta = \mu_0\} \in J_n$*)*
- (γ) *if* $0 < n < \omega, \alpha < \kappa_n$ *and* $\eta \in \prod\limits_{\ell<n} \kappa_\ell$ *then* $\lambda_\eta > \mu_0 \Rightarrow \lambda_\eta > \lambda_{\eta\hat{\ }\langle\alpha\rangle}$ *so* $\{\eta \in \prod\limits_{\ell<n} \kappa_\ell : \lambda_{\eta\hat{\ }\langle\alpha\rangle} \nleq \lambda_\eta\} \in J_n$ *hence*
- $(\gamma)'$ $\{\eta \in \prod\limits_{\ell\le n} \kappa_\ell : \lambda_\eta \nleq \lambda_{\eta\restriction n}\} \in J_{n+1}$
- (δ) $J_n = \{A \subseteq \prod\limits_{\ell<n} \kappa_\ell :$ *max pcf*$\{\lambda_\eta : \eta \in A\} < \mu\}$.

Question: Can we prepare the ground to 3.8 with IND$^+$ instead IND?

Proof. We choose $\bar{\lambda}^n$ by induction on n. For $n = 1$ apply [Sh:g, Ch.II,1.5A] to the sequence $\langle \lambda^1_i : i < \kappa_0\rangle$, the ideal $\{A \subseteq \kappa_1 : \max \mathrm{pcf}\{\lambda^1_i : i < \kappa_0\} < \mu_0\}$ and the cardinal μ and get $\langle \lambda_{<i>} : i < \kappa_0\rangle$. For $n+1$ for each $i < \lambda_n$ we apply [Sh:g, Ch.II,1.5A] to $\langle \lambda_n : \eta \in \prod\limits_{\ell<n} \kappa_\ell\rangle$, the ideal $\{A \subseteq \prod\limits_{\ell<n} \kappa_\ell : \max \mathrm{pcf}\{\lambda_\eta : \eta \in A\} < \mu\}$ and the cardinal λ^n_i and we get $\langle \lambda_{\eta\hat{\ }<i>} : \eta \in \prod\limits_{\ell<n} \kappa_\ell\rangle$. $\square_{3.4}$

3.5 Claim. *In 3.4, from* $\otimes_2$ *we can deduce*

> $\otimes^3$ *there are functions* $f_{\ell,n} : \prod\limits_{m=\ell+1}^{n} \kappa_m \rightarrow I_\ell$ *(for* $\ell < N < \omega$*) such that for every* $\eta \in \prod\limits_{m<\omega} \kappa_m$ *for some* $\ell < n < \omega$ *we have* $\eta(\ell) \in f\ell, n(\eta \restriction (\ell, n))$.

Proof. Otherwise $\langle \lambda_{\eta\restriction n} : n < \omega\rangle$ is a strictly decreasing sequence of cardinals. $\square_{3.5}$

3.6 Definition. 1) IND$(\langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle)$ (note that $|\mathrm{Dom}(J_\varepsilon)|$ is not necessarily increasing with ε) means that eac J_ε is an ideal on $\mathrm{Dom}(J_\varepsilon)$, say κ_ε and

- $(*)$ for every sequence $\langle f_{\varepsilon,u} : \varepsilon < \varepsilon^*, u \subseteq \varepsilon^* \backslash (\varepsilon+1)$ finite$\rangle$ such that $f_{\varepsilon,u}$ a function from $\prod\limits_{\zeta\in u} \kappa_\zeta$ to J_ε there is an increasing sequence $\varepsilon_0 < \varepsilon_1 < \ldots < \varepsilon_n < \ldots < \varepsilon^*$ (for $n < \omega$) and $\alpha_\ell \in \kappa_{\varepsilon_\ell}$ (for $\ell < \omega$) such that:
- $(**)$ for $\ell < n < \omega$ we have $\alpha_\ell \notin f_{\varepsilon_\ell,u}(\langle \alpha_{\varepsilon_{\ell+1}}, \ldots, \alpha_{\varepsilon_n}\rangle)$ for $u = \{\alpha_{\varepsilon_{\ell+1}}, \ldots, \alpha_{\varepsilon_n}\}$.

2) $\mathrm{IND}^+(\langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle)$ means J_ε is an ideal (on $\mathrm{Dom}(J_\varepsilon)$ which is say κ_ε) such that:

$(*)$ for every sequence $\langle f_{\varepsilon,u} : \varepsilon < \varepsilon^*, u \subseteq \varepsilon^* \backslash (\varepsilon+1) \text{ finite}\rangle$ such that $f_{\varepsilon,u} : \prod\limits_{\zeta \in u} \kappa_\zeta \to J_\varepsilon$ there is $\langle \alpha_\varepsilon : \varepsilon < \varepsilon^*\rangle \in \Pi\kappa_\varepsilon$ such that

$(**)$ for $\varepsilon < \varepsilon^*, u \subseteq \varepsilon^* \backslash (\varepsilon+1)$ finite we have: $\alpha_\varepsilon \notin f_{\varepsilon,u}(\dots, \alpha_\zeta, \dots)_{\zeta \in u}$.

3) Let function $\langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle$ be the set of $\bar f$ as in $(*)$ of part (1), i.e. $\bar f = \langle_{\varepsilon,u}: \varepsilon < \varepsilon^*$ and $u \subseteq \varepsilon^* \backslash (\varepsilon+1)$ is finite$\rangle$ where $f_{\varepsilon,u} \in \prod\limits_{\zeta \in u} \kappa_\zeta$. We say for $\bar f \in$ function$\langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle$, that $\bar\varepsilon$ is candidate if $\bar\varepsilon$ is an increasing sequence of length ω of ordinals $< \varepsilon^*$. In this case we say that $\bar\alpha$ is $(\bar f, \bar\varepsilon)$-free if $\bar\alpha \in \prod\limits_{n,\omega} \mathrm{Dom}(J_{\varepsilon_n})$ and the statement $(**)$ of part (1) holds.

3.7 Observation. 1) If $\mathrm{IND}(\bar J)$ where $\bar J = \langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle$ is as in 3.6, each J_ε is $(|\varepsilon^*|^{\aleph_0})^+$-complete, then for some $\varepsilon_0 < \varepsilon_1 < \ldots < \varepsilon^*$ we have $\mathrm{IND}^+(\langle J_{\varepsilon_n} : n < \omega\rangle)$.
2) If $\mathrm{IND}(\bar J)$ where $\bar J = \langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle$ as in 3.6, each J_ε is $\mathrm{cov}(|\varepsilon^*|, \mu, \aleph_1, 2)^+$-complete then for some infinite $u \in \mathscr{S}_{<\mu}(\varepsilon^*)$ we have $\mathrm{IND}(\bar J \restriction u)$.

Before we prove 3.7

3.8 Conjecture. if $\mathrm{IND}(\langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle), \varepsilon^* < \omega_1$ and each J_ε is $\aleph_1$-complete then for some c.c.c. forcing P we have:

$$\Vdash_P \text{“ for some } \varepsilon_0 < \ldots < \varepsilon_n < \varepsilon_{n+1} < \ldots < \varepsilon^*,\ \mathrm{IND}(\langle J_{\varepsilon_n} : n < \omega\rangle)\text{”}.$$

3.9 Remark. In the proof of 3.7(2) it is enough to demand on $\mathscr{P}$:

$(*)$ if $\varepsilon_0 < \varepsilon_1 < \ldots < \varepsilon_n < \ldots < \varepsilon^*$ (for $n < \omega$) then for some $b \in \mathscr{P}$, $(\exists^\infty n)\varepsilon_n \in b$

this seems to weaken $\mathrm{cov}(\dots)$ but does not.

Proof of 3.7. 1) Similar to the proof of part (2), as $\mathrm{cov}(\lambda, \aleph_1, \aleph_2, 2) \le |\varepsilon^*|^{\aleph_0}$.
2) Let $\mu =:\ \mathrm{cov}|\varepsilon^*, \mu, \aleph_1, 2)$ and let $\mathscr{P} \subseteq [\varepsilon^*]^{<\mu}$ be of cardinality μ exemplifying its definition i.e. $(\forall a)[a \subseteq \varepsilon^*\ \&\ |a| = \aleph_0 \Rightarrow (\exists b \in \mathscr{P})[a \subseteq b]]$.

If for some $b \in \mathscr{P}$, $\mathrm{IND}(\bar J \restriction u)$ hold then we are done. Otherwise for each $b \in \mathscr{P}$, we can find $\bar f^b = \langle f^b_{\varepsilon,u} : \varepsilon \in b \text{ and } b \subseteq u \backslash (\varepsilon+1) \text{ is finite}\rangle \in$ function$(J \restriction u)$ such that for no $\bar\varepsilon = \langle \varepsilon_n : n < \omega\rangle$ strictly increasing sequence of ordinals from b and $\alpha_n \in \mathrm{Dom}(J_{\varepsilon_n})$ (for $n < \omega$) do we have $n < \omega\ \&\ u \subseteq \{\varepsilon_{n+1}, \varepsilon_{n+2}, \dots\} \Rightarrow \alpha_n \notin f^b_{\varepsilon,u}(\{\alpha_m : m \in u\})$. Let us define for $\varepsilon < \varepsilon^*$ and $u \subseteq \varepsilon^* \backslash (\varepsilon+1)$ finite a function $f_{u,\varepsilon}$ from $\prod\limits_{\zeta \in u} \mathrm{Dom}(J_\zeta)$ to J_ε by:

$(*)$ if $\alpha_\zeta \in \mathrm{Dom}(J_\zeta)$ for $\zeta \in u$ then $f_{\zeta,u}(\dots, \alpha_\zeta, \dots)_{\zeta \in u} = \bigcup\{f^b_{\zeta,u} : \zeta \in b \text{ and } u \subseteq b \text{ and } b \in \mathscr{P}\}$.

(As each J_ζ is μ^+-complete (by assumption) $\text{Rang}(f_{\zeta,u}) \subseteq J_\zeta$. As $\text{IND}(\langle J_\varepsilon : \varepsilon < \varepsilon^* \rangle)$ necessarily there is a strictly increasing $\langle \varepsilon_n : n < \omega \rangle, \varepsilon_n < \varepsilon^*$, and $\alpha_n \in \text{Dom}(J_{\varepsilon_n})$ (for $n < \omega$) such that:

$(**)$ $n < \omega, u \subset \{\varepsilon_{n+1}, \varepsilon_{n+2}, \dots\}$ finite, implies $\alpha_n \notin f_{\varepsilon_n,u}(\dots, \alpha_m, \dots)_{\varepsilon_m \in u}$.

By the choice of $\mathscr{P}$ for some $b \in \mathscr{P}$ we have $\{\varepsilon_n : n < \omega\} \subseteq b$, but then $\langle (\varepsilon_n, \alpha_n) : n < \omega \rangle$ contradict the choice of $\bar f^b = \langle f^b_{\zeta,u} : \zeta \in b, u \subseteq b \backslash (\zeta+1) \text{ finite} \rangle$. $\square_{3.7}$

3.10 Conclusion. 1) If $\text{IND}^+(\langle I_n : n < \omega \rangle)$ and $\text{Dom}(I_n) = \kappa_n$ then

- (a) the conclusion of 3.5 (i.e. $\otimes^3$ there) fails, hence $\otimes^2$ of 3.4 fails hence $\otimes^1$ of 3.4 fails
- (b) if $\lambda > \sum\limits_{n<\omega} \kappa_n, \kappa_n < \text{cf}(\kappa_{n+1})$, then for every n large enough for no $\lambda_i \in (\sum\limits_{n<\omega} \kappa_n, \lambda) \cap \text{Reg}$ (for $i < \kappa_n$) is $\prod\limits_{i<\kappa_n} \lambda_i / I_n \lambda$-directed.

2) If we weaken the assumption to $\text{IND}(\langle I_n : n < \omega \rangle)$ then in (b) we have just for arbitrarily large $n < \omega$.

Proof. 1) ?

- (a) straight
- (b) our problem is to get μ_1, which is not serious.

2) Similarly. $\square_{3.10}$

3.11 Conclusion. 1) Assume $\langle \kappa_\varepsilon : \varepsilon < \delta \rangle$ is strictly increasing, $|\delta| \le \sigma < \kappa_0, \kappa = \sum\limits_{i<\delta} \kappa_i$ and $\text{IND}(\kappa, \sigma)$. If $\lambda > \kappa$ then for every large enough $\varepsilon < \delta$, there are no $\lambda_\alpha \in (\kappa, \lambda) \cap \text{Reg}$ for $\alpha < \kappa_\varepsilon$ such that $\prod\limits_{\alpha<\kappa_\varepsilon} \lambda_\alpha / [\kappa_\varepsilon]^{\le\sigma}$ is λ-directed recalling $[\kappa_\varepsilon]^{\le\sigma} = \{a \subseteq \kappa_\varepsilon : |a| \le \sigma\}$.
2) If $\text{IND}(\kappa)$, $\text{cf}(\kappa) = \aleph_0 = \sigma, \kappa = \sum\limits_{n<\omega} \kappa_n, \kappa_n < \kappa_{n+1}$ then the conclusion of 1) holds.
3) If $\text{IND}(\kappa, \sigma), \delta = \omega, \kappa_\varepsilon = \kappa$, then the conclusion of (1) holds.
4) If $\varepsilon < \lambda_0, \langle \lambda_\varepsilon : \varepsilon < \varepsilon^* \rangle$ is a strictly increasing sequence of regular cardinals, J_ε is an ideal on λ_ε extending $J^{\text{bd}}_{\lambda_\varepsilon}$ and we have $\text{IND}(\langle J_\varepsilon : \varepsilon < \varepsilon^* \rangle)$ then for some $\varepsilon' \le \varepsilon^*$, $\text{cf}(\varepsilon') = \aleph_0$ and not only $\text{IND}(\langle J_\varepsilon : \varepsilon' \rangle)$ but we can even demand $\varepsilon' = \bigcup\limits_{n<\omega} \varepsilon_n$.
5) $\text{IND}(\langle \lambda_n : n < \omega \rangle)$ iff $\text{IND}^+(\langle [\lambda_n]^{\le \lambda_{n-1}} : n < \omega \rangle)$ (stipulating $\lambda_{-1} = \aleph_0$).
6) If $\text{IND}^0(\lambda, \sigma)$ then $\text{IND}^+(\langle \lambda : n < \omega \rangle)$.

Proof. Check.

3.12 Discussion: Let $\bar J = \langle J_\varepsilon : \varepsilon < \varepsilon^* \rangle$ and assume $\text{IND}(\bar J)$.
1) Note that if P is a θ-c.c. forcing notion, each J_ε is θ-complete then for any

$\bar f = \langle f_{\varepsilon,u} : \varepsilon < \varepsilon^*, u \in [\varepsilon^* \backslash (\varepsilon+1)]^{<\aleph_0}\rangle \in V^P$ as in Definition 3.6 we can find $\bar f' = \langle f'_{\varepsilon,u} : \varepsilon < \varepsilon^*, u \in [\varepsilon^* \backslash (\varepsilon+1)]^{<\aleph_0}\rangle \in V$ such that for every $\bar\alpha \in \prod\limits_{\zeta \in u} \mathrm{Dom}(J_\zeta)$ we have $f_{\varepsilon,u}(\bar\alpha) \subseteq f'_{\varepsilon,u}(\bar\lambda)$, so we can consider only $\bar f \in V$. For each such $\bar f$ let $A_{\bar f} = \{v :$ for some strictly increasing sequence $\bar\varepsilon = \langle \varepsilon_n : n < \omega\rangle$ of ordinals $< \varepsilon^*$ and $\bar\alpha \in \prod\limits_{n<\omega} \mathrm{Dom}(J_{\varepsilon_n})$ the conclusion $(**)$ of Definition 3.6 holds and $v = \{\varepsilon_\ell : \ell < m\}$ for some $m < \omega\}$.

For $\bar g, \bar f \in$ function$(\bar J)$ where $\bar J = \langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle$, let $\bar g \le \bar f$ iff for every $\varepsilon < \varepsilon^*$ and $u \in [\varepsilon^* \backslash (\varepsilon+1)]^{<\aleph_0}$ we have $\bar\alpha \in \prod\limits_{\varepsilon \in u} \mathrm{Dom}(J_\varepsilon) \Rightarrow g_{\varepsilon,u}(\bar\alpha) \subseteq f_{\varepsilon,u}(\bar\alpha)$. Clearly $\bar g \le \bar f \Rightarrow A_{\bar f} \subseteq A_g$.

2) In V we can define a filter D on $[\bigcup\limits_{\varepsilon<\varepsilon^*} \mathrm{Dom}(\bar J_\varepsilon)]^{<\aleph_0}$:

$$A \in D \ \underline{\text{iff}} \text{ for some } \bar f \in \text{ function}(\langle J_\varepsilon : \varepsilon < \varepsilon^*\rangle) \text{ we have } A_f \subseteq A.$$

Now $D \subseteq \mathscr{P}([\bigcup\limits_{\varepsilon<\varepsilon^*} \mathrm{Dom}(J_\varepsilon)]^{<\aleph_0})$ and D is upward closed trivially. Also D is closed under intersection of countable many members if each J_n is $\aleph_1$-closed (similarly σ-closed) because if $A_n \in D$ let $\bar f^n \in$ function$(\bar J)$ be such that $A_{\bar f^n} \subseteq A$. Now for some $g \in$ function$(J)[n < \omega \Rightarrow \bar f^n \subseteq \bar g]$, hence $A_g \subseteq A_{\bar f^n}$, so $A_{\bar g} \subseteq A_n$ for $n < \omega$ and obviously $A_{\bar g} \in D$. Lastly $\emptyset \notin D$ as $\mathrm{IND}(\bar J)$ holds.

3.13 Claim. *Suppose for $\alpha < \alpha^*, \bar I^\alpha = \langle I^\alpha_n : n < \omega\rangle, \kappa = \sup\{|Dom(I^\alpha_n)| : \alpha < \alpha^*\}$, $IND^+(\langle I^\alpha_n : n < \omega\rangle)$, and:*

- $(*)$ *if $\alpha < \alpha^*, f_n : Dom(I^\alpha_n) \to$ Ord* *__then__* *for some $n(*) < \omega, \beta < \alpha^*$ and ordinal γ*

$$I^\beta_{1+n} \cong I^\alpha_{n(*)+1+n} \restriction \{x \in Dom(I^\alpha_{n(*)+1+n}) : f_{1+n}(x) > \gamma\}$$

$$I^\beta_0 = I^\alpha_{n(*)} \restriction \{x \in Dom(I^\alpha_{n(*)}) : f_0(x) < \gamma\}.$$

__Then__ for no $\lambda > \kappa$ and $\alpha < \alpha^*$ do we have for every $n < \omega$:

$$x \in Dom(I^\alpha_n) \Rightarrow \lambda^n_x \in (\kappa, \lambda) \cap Reg \text{ and } \prod_{x \in \mathrm{Dom}(I^\alpha_n)} \lambda^n_x / I^\alpha_n \text{ is } \lambda^+\text{-directed.}$$

Proof. No new point, like the proof of 2.6.

Remark. 1) This claim is used in the proof of 5.4.
2) If in 3.13, $\alpha^* = 1$ and I^α_{n+1} is λ_n-complete, $\lambda_n > |\mathrm{Dom}(I^\alpha_\ell))|$ for $\ell < n$ then $(*)$ there holds.

__Question__: If $\mathrm{IND}(\lambda,\sigma)$, $\mathrm{cf}(\lambda) = \aleph_0$ do we have $\mathrm{IND}(\langle J^{\mathrm{bd}}_{\lambda_n} : n < \omega\rangle)$ for some $\lambda_n = \mathrm{cf}(\lambda_n) < \lambda, \sigma < \lambda_n$?

3.14 Claim. *In 2.6(1) we can deduce even IND*($\langle J^{\rm bd}_\theta : \theta \in \mathfrak{a}\rangle$).

Proof. Reread the proof of 2.6: let $f_{u,\varepsilon}$ be as in Definition 3.6, so without loss of generality $\text{Rang}(f_{\varepsilon,u}) \subseteq \kappa_\varepsilon$ (as $\alpha = \{\beta : \beta < \alpha\}$) and $M = (\kappa, \kappa_\varepsilon, f_{u,\varepsilon})_{\varepsilon,u}$. Now repeat the proof of 2.6. $\square_{3.14}$

Next we improve the ideals from "bounded" to "nonstationary"

3.15 Claim. *1) Assume* $\lambda > cf(\lambda), |\mathfrak{a}| < min(\mathfrak{a})$ *and* $\lambda = \sup(\mathfrak{a})$ *and* $rk'_{\langle J_{<\theta}[\mathfrak{a}]:\theta\in\text{pcf}(\mathfrak{a})\rangle}(\mathfrak{a}) \ge |\mathfrak{a}|^+, \sigma^* \in (|\mathfrak{a}|, min(\mathfrak{a})) \cap$ *Reg, and*

$$I_\theta =: \{S : S \subseteq \theta \textit{ and } \{\delta \in S : cf(\delta) = \sigma^*\} \textit{ is not stationary}\}$$

then IND($\langle I_\theta : \theta \in \mathfrak{a}\rangle$).
2) Moreover for any sequence $\bar H = \langle H_\theta : \theta \in \mathfrak{a}\rangle$, *where* $\mathfrak{a} \subseteq \lambda, H_\theta$ *a function from* $[\lambda]^{<\aleph_0}$ *to* I_θ *and* $\mathfrak{c}, \mathfrak{a} \subseteq \mathfrak{c} \subseteq pcf(\mathfrak{a}), |\mathfrak{c}| < min(\mathfrak{a})$ *we can find* $\bar\alpha = \langle \alpha_\tau : \tau \in \mathfrak{a}\rangle \in \Pi\mathfrak{a}$ *such that defining for* $\mathfrak{b} \subseteq \mathfrak{a}$

$$c\ell_{\bar H,\bar\alpha}(\mathfrak{b}) = \{\tau \in \mathfrak{a}, \alpha_\tau \in H_\tau(\bar\alpha \restriction \mathfrak{e}) \textit{ for some finite } \mathfrak{e} \subseteq \mathfrak{b}\}$$

we have

$$(*) \quad \theta \in \mathfrak{c} \;\&\; \mathfrak{b} \in J_{<\theta}[\mathfrak{a}] \Rightarrow c\ell_{\bar H,\bar\alpha}(\mathfrak{b}) \in J_{<\theta}[\mathfrak{a}].$$

Proof. 1) We can prove it from part 2) exactly as in the proof of 2.6(1).
2) Let $\bar{\mathfrak{b}} = \langle \mathfrak{b}_\theta[\mathfrak{a}] : \theta \in \text{pcf}(\mathfrak{a})\rangle$ be a generating sequence for $\mathfrak{a}$, without loss of generality $\max \text{pcf}(\mathfrak{a}) \in \mathfrak{c}$ and

$$[\theta \in \mathfrak{c} \backslash \max \text{pcf}(\mathfrak{a}) \Rightarrow \min(\text{pcf}(\mathfrak{a}) \backslash \theta^+) \in \mathfrak{c}].$$

Before we continue, recall that we know:
3.16 Fact If $|\mathfrak{a}| < \min(\mathfrak{a}), \mathfrak{a} \subseteq$ Reg and $\langle \mathfrak{b}_\sigma[\mathfrak{a}] : \sigma \in \text{pcf}(\mathfrak{a})\rangle$ a generating sequence for $\mathfrak{a}$ then

$(*)_\mathfrak{a}$ there is $\langle \bar f^\theta : \theta \in \text{pcf}(\mathfrak{a})\rangle$ such that
- (a) $\bar f^\theta = \langle f^\theta_\alpha : \alpha < \theta\rangle$ is $<_{J_{<\theta}[\mathfrak{a}]}$-increasing
- (b) $\bar f^\theta$ is cofinal in $(\Pi\mathfrak{a}, <_{J_\theta[\mathfrak{a}]})$ where $J_\theta[\mathfrak{a}] =: J_{<\theta}[\mathfrak{a}] + (\mathfrak{a} \backslash \mathfrak{b}_\theta[\mathfrak{a}])$
- (c) if $\sigma \in \theta \cap \mathfrak{c}, \mathfrak{a} < \theta$ and $\mathfrak{b} = \mathfrak{b}_\sigma[\mathfrak{a}]$ then for some $n < \omega$ and $\alpha_\ell < \theta_\ell \le \sigma$ (for $\ell \le n$) we have $f^\theta_\delta \restriction \mathfrak{b} = (\max\{f^{\theta_\ell}_{\alpha_\ell} : \ell < n\}) \restriction \mathfrak{b}$ (the max is pointwise)
- (d) if $\delta < \theta \in \text{pcf}(\mathfrak{a})$, $\text{cf}(\delta) \in (|\mathfrak{a}|, \min(\mathfrak{a}))$ then for every $\tau \in \mathfrak{a}$

$$f^\theta_\delta(\tau) = \min\{\cup\{f^\theta_\alpha(\tau) : \alpha \in C\} : C \text{ a club of } \delta\}$$

provided that the function defined satisfies condition (c) above

(exist by [Sh:g, Ch.VIII,§1] or [Sh 430, 6.7x] we choose by induction on θ).

Let

$$\begin{aligned} S^{\rm gd}_\theta =: \big\{\delta < \theta : \mathrm{cf}(\delta) \in (|\mathfrak{a}|, \min(\mathfrak{a})), f^\theta_\delta \text{ is a } <_{J^{\rm bd}_{<\theta}[\mathfrak{a}]}\text{-eub of } \bar{f}^\theta \restriction \delta \text{ and} \\ \{\tau \in \mathfrak{a} : \mathrm{cf}(f^\theta_\alpha(\tau)) = \mathrm{cf}(\delta)\} = \mathfrak{b}_\theta[\mathfrak{a}] \bmod J_{<\theta}[\mathfrak{a}]\big\} \end{aligned}$$

(alternatively use simultaneous witnesses for $I[\theta]$ as in [Sh 420, §1].
Note:

$(*)_{\mathfrak{a}}$ (e) if E_τ is a club of τ for $\tau \in \mathfrak{a}$ and $\theta \in \mathrm{pcf}(\mathfrak{a})$ then for some club E of θ:

$$\delta \in E \cap S^{\rm gd}_\theta \Rightarrow \{\tau \in \mathfrak{a} : f^\theta_\delta(\tau) \in E_\tau\} = \mathfrak{a} \bmod J_\theta[\mathfrak{a}].$$

Let $\bar{H} = \langle H_\tau : \tau \in \mathfrak{a}\rangle$ be as in the claim, so H_τ is a function from $[\lambda]^{<\aleph_0}$ to I_τ. Now for every $\theta \in \mathrm{pcf}(\mathfrak{a})$ and $\alpha < \theta$ and $\tau \in \mathfrak{a}$ we define $A^{\theta,\tau}_\alpha = A^\tau(f^\theta_\alpha) \in I_\tau$ as

$$\bigcup \{H_\tau(u) : u \subseteq \mathrm{Rang}(f^\theta_\alpha) \text{ is finite}\}$$

(as I_τ is τ-complete, $\tau > |\mathfrak{a}| = |\{u : u \subseteq \mathrm{Rang}(f^\theta_\alpha) \text{ finite}\}|$, really $A^\theta_\alpha \in I_\tau$).

Now for each $\alpha < \theta \in \mathrm{pcf}(\mathfrak{a})$ and $\sigma \in \mathrm{pcf}(\mathfrak{a})$ by $(*)_{\mathfrak{a}}(e)$ applied with $\sigma, \langle A^{\theta,\tau}_\alpha : \tau \in \mathfrak{a}\rangle$ here standing for $\theta, \langle E_\tau : \tau \in \mathfrak{a}\rangle$ there, we get a club $C_{\theta,\sigma,\alpha}$ of σ.

For each $\sigma \in \mathrm{pcf}(\mathfrak{a})$ let $C_\sigma = \bigcap\limits_{\theta \in \mathfrak{c} \cap \sigma, \alpha < \theta} C_{\theta,\sigma,\alpha}$ (note: $|\mathfrak{c} \cap \sigma| < \sigma$), so C_σ is a club of σ. Lastly let $\sigma^* = \mathrm{cf}(\sigma^*) \in (|\mathfrak{a}|, \min(\mathfrak{a}))$ and $\langle N_i : i \le \sigma^*\rangle$ be as in [Sh:g, Ch.VIII,1.2,1.4], so

$\oplus_1$ $N_i \prec (\mathscr{H}(\chi), \in, <^*_\chi)$ is increasing continuous, for $i \le \sigma^*, \|N_i\| = \sigma^*$, $\langle N_j : j \le i\rangle \in N_{i+1}$.

Let $\alpha_\theta =: \sup(N_{\sigma^*} \cap \theta)$ for $\theta \in \mathfrak{c}$; so $\alpha_\theta \in S^{\rm gd}_\theta$ for every $\theta \in \mathfrak{c}$, and we shall show that $\bar{\alpha} = \langle \alpha_\tau : \tau \in \mathfrak{a}\rangle$ is as required.

For each $\sigma, \theta \in \mathfrak{c}, \sigma < \theta$ we have: $\alpha_\theta \in C_\theta$ hence $\alpha_\theta \in C_{\theta,\sigma,\alpha_\sigma}$ hence

$$\{\tau \in \mathfrak{a} : f^\sigma_{\alpha_\sigma}(\tau) \in \tau \backslash A^{\sigma,\tau}_{\alpha_\sigma}\} = \mathfrak{b}_\tau[\mathfrak{a}] \bmod J_\sigma[\mathfrak{a}].$$

The rest should be clear. $\square_{3.15}$

We next point out another connection; if the rank is small and $|\mathrm{pcf}(\mathfrak{a})|$ is large, then we have a case of "$\Pi\mathfrak{d}/\mathscr{S}_{\le\lambda}(\mathfrak{d})$ has large true cofinality".

3.17 Claim. *If $\zeta > \mathrm{rk}'(\mathfrak{a}, \langle J_{<\theta}[\mathfrak{a}] : \theta \in \mathrm{pcf}(\mathfrak{a})\rangle)$ and $\bar{\lambda} = \langle \lambda_\varepsilon : \varepsilon \le \zeta\rangle$ is strictly increasing and $|\mathrm{pcf}(\mathfrak{a})| \ge \lambda_\zeta$ then for some $\varepsilon < \zeta$ and $\mathfrak{c} \subseteq \mathfrak{a}$ we have $|\mathrm{pcf}(\mathfrak{c})| \ge \lambda_{\varepsilon+1}$ and $\mathfrak{d} \in J_*[\mathrm{pcf}(\mathfrak{c}) \Rightarrow \Pi\mathfrak{d}/\mathscr{S}_{\le\lambda_\varepsilon}(\mathfrak{d})$ has the true cofinality which is $\max \mathrm{pcf}(\mathfrak{c})$.*

Proof. If not, prove by induction on $\varepsilon \le \zeta$ that

$(*)$ if $\mathfrak{c} \subseteq \mathfrak{a}, \varepsilon = \mathrm{rk}'(\mathfrak{c}, \langle J_{<\theta}[\mathfrak{a}] : \theta \in \mathrm{pcf}(\mathfrak{a})\rangle)$ then $|\mathrm{pcf}(\mathfrak{c})| < \lambda_{\varepsilon+1}$.

Let $J_0 = \{\mathfrak{d} : \mathfrak{d} \subseteq \mathfrak{c}, \varepsilon > \text{ rk}'(\mathfrak{d}, \langle J_{<\theta}[\mathfrak{a}] : \theta \in \text{ pcf}(\mathfrak{a})\rangle)\}$. By the induction hypothesis $[\mathfrak{d} \in J_0 \Rightarrow |\text{pcf}(\mathfrak{d}) \le \lambda_\varepsilon]$. Let J be the ideal on $\mathfrak{c}$ which J_0 generates. We have: $[\mathfrak{d} \in J \Rightarrow |\text{pcf}(\mathfrak{d})| \le \lambda_\varepsilon]$, so by the assumption toward contradiction $\mathfrak{c} \notin J$. Let $\theta = \max \text{pcf}(\mathfrak{c})$. So by the definition of $\text{rk}' : J_{<\theta}[\mathfrak{c}] \subseteq J$. Hence (see [Sh:g, Ch.VIII,§3] or [Sh:E11] for $\mathfrak{d} \subseteq \text{ pcf}(\mathfrak{c}), \mathfrak{d} \in J_*[\text{pcf}(\mathfrak{a})]$ we have $\Pi\mathfrak{d}/J^*_{<\theta}[\text{pcf}(\mathfrak{c})]$ has the true cofinality θ and $J_*[\text{pcf}(\mathfrak{a})]$ is generated by $\{\text{pcf}(\mathfrak{b}) : \mathfrak{b} \in J_{<\theta}[\mathfrak{c}]\}$. So the conclusion holds. $\square_{3.17}$

§4 Sticks and BA's

4.1 Lemma. *Assume* $\theta \leq \mu < \lambda \leq \lambda^*, J$ *an ideal on* θ *and assume*

$\otimes^J_{\theta,\mu,\lambda,\lambda^*}$ *if* $n < \omega, \mathfrak{a}_i \in [\mathit{Rang}\ \cap \lambda^+ \backslash \mu^+]^n$ *for* $i < \theta$ *then*
$\{a \in J : \max \mathit{pcf}(\bigcup\limits_{i \in a} \mathfrak{a}_i) \leq \lambda^*\}$ *is generated by* $\leq \mu$ *sets.*[1]

Then there is a set H *such that*

- (a) H *a set of partial functions from* θ *to* $[\lambda]^{\leq \mu}$
- (b) $|H| \leq \lambda^*$
- (c) *for every function* $g : \theta \rightarrow \lambda$ *we can find* h *and* $\bar{\mathfrak{a}} = \langle \mathfrak{a}_i : i < \theta \rangle$ *such that*
  - (i) $\mathfrak{a}_i$ *is a finite set of regular cardinals from* $(\theta, \lambda]$
  - (ii) h *is a function from* θ *to* $[\lambda]^{\leq \mu}$ *such that* $\bigwedge\limits_i g(i) \in h(i)$
  - (iii) *for any* $n < \omega$ *and* $a \subseteq \theta$*:*
    if $(\forall i \in a)[|\mathfrak{a}_i| \leq n]$ *and max pcf*$(\bigcup\limits_{i \in a} \mathfrak{a}_i) \leq \lambda^*$ *then for some* b *satisfying*
    $a \subseteq b \subseteq \theta$ *we have* $h \restriction b \in H$.

Proof. Like [Sh 430, §2].

4.2 Remark. 1) But we can then change the bound (in clause (c)(ii)) to $h(i) \in [\lambda]^{<\mu}$. Then $\otimes_{\theta,\mu,\lambda,\lambda^*}$ is changed to

$\otimes'_{\theta,\mu,\lambda,\lambda^*}$ if $n < \omega, \mathfrak{a}_i \in [\mathrm{Reg}\ \cap \lambda^+]^n$ for $i < \theta$ then
$\{a \in J : \text{for some } \mu_0 < \mu, \max \mathrm{pcf}(\bigcup\limits_{i \in a} \mathfrak{a}_i \backslash \mu_0^+) \leq \lambda^*\}$
is generated by $< \mu$ sets.

2) We can weaken $\otimes$ to

$\otimes^J_{\theta,\mu,\lambda,\lambda^*}$ if $n < \omega, \mathfrak{a}_i \in [\mathrm{Reg}\ \cap \lambda^+]^n$ for $i < \theta$ then
$\{a \in J : \max \mathrm{pcf}(\bigcup\limits_{i < a} \mathfrak{a}_i) \leq \lambda^*\}$
is generated (as an ideal) by some $\mathscr{P} \subseteq J$ such that

$$\kappa \in \bigcup_{i < \theta} \mathfrak{a}_i \Rightarrow \kappa < \{a \in \mathscr{P} : \kappa \in \bigvee_{i \in a} \mathfrak{a}_i\}.$$

---

[1] see 4.2(3)

3) Instead $\otimes_{\theta,\mu,\lambda,\lambda^*}$ we can define a game

$\otimes_{\theta,\mu,\lambda,\lambda^*}[\mathfrak{D}]$ First player has no winning strategy in the game defined below

$GM'_{\theta,\mu,\lambda,\lambda^*}[\mathfrak{D}]$ The play lasts ω moves, in the n-th move:
first player chooses $\bar{\lambda}^n = \langle \lambda^n_i : i \in A_n \rangle, A_0 = \theta, \bigwedge\limits_{m<n} A_m \subseteq A_n, \bigwedge\limits_{m<n} \bigwedge\limits_{i \in A_n} \lambda^n_i < \lambda^m_i, \lambda^n_i = \text{cf}(\lambda^n_i) \in (\mu, \lambda]$ and
second player chooses an ideal J_n on $A_n, J_n \subseteq \{a \subseteq A_n : \max \text{pcf}\{\lambda^n_i : i \in a\} \le \lambda^*\}, J_n$ generated by $\le \theta$ sets.
In the end (clearly $\bigcap\limits_{n<\omega} A_n = \emptyset$) they produce the ideal J, the one generated by $\{a \subseteq \theta : \text{for some } n, a \subseteq A_n \backslash A_{n+1} \text{ and } a \in J_n\}$.
Second player wins if $J \in \mathfrak{D}$.

4.3 Definition. Assume $\bar{J} = \langle J_\ell : \ell < 3 \rangle$, where $J_0 \subseteq J_1 \subseteq J_2 \subseteq \mathscr{P}(\theta)$, each J_ℓ is downward closed (usually is an ideal); we let $J^+_\ell =: \mathscr{P}(\theta) \backslash J_\ell$.

1) $\text{dcf}_{\bar{J}}(\lambda < \mu) = \min\Big\{|\mathscr{F}| : \mathscr{F}$ is a family of functions each with domain from J^+_1 and range included in $[\lambda]^{<\mu}$ such that:
$(*)_{\mathscr{F}}$ for every $b \in J^+_2$ and $f \in {}^b\lambda$ for some $a \in J^+_1$ and $g \in \mathscr{F} \cap {}^a\lambda$ we have $(\forall^{J_0} i \in a)(i \in b \ \& \ g(i) \in f(i))$
i.e. $\{i : i \in a \text{ and } i \notin b \vee g(i) \notin f(i)\} \in J_0\Big\}$

2) $\text{ecf}_{\bar{J}}(\lambda < \mu) = \min\Big\{|\mathscr{F}| : \mathscr{F}$ is a family of functions each with domain from J^+_1 and range included in $[\lambda]^{<\mu}$ such that:
$(*)_{\mathscr{F}}$ for every $b \in J^+_2$ and $f \in {}^b\lambda$ for some $a \in S^+_1$ and $g \in \mathscr{F} \cap {}^\alpha\lambda$ we have
$(\exists^{J_0} i \in a)(i \in b \ \& \ g(i) \in f(i))$
i.e. $\{i : i \in a, i \in b \text{ and } g(i) \in f(i)\} \notin J_0\Big\}$

3) $\text{xcf}_{\bar{J}}(\lambda, < \mu^+)$ is written $\text{xcf}_{\bar{J}}(\lambda, \le \mu)$ or $\text{xcf}_{\bar{J}}(\lambda, \mu)$. Also $\text{xcf}_{\bar{J}}(\lambda, 1)$ is written $\text{xcf}_{\bar{J}}(\lambda)$. Also $\text{xcf}_{<[\lambda]^{<\sigma},[\lambda]^{<\theta},[\lambda]^{<\chi}}(\lambda, < \mu)$ is written $\text{xcf}_{\chi,\theta,\sigma}(\lambda, < \mu)$, etc.

4.4 Definition. $\text{⫯}_{\lambda,\mu,\theta} =: \min\{|\mathscr{P}| : \mathscr{P} \subseteq [\lambda]^\theta$ is such that for every $A \in [\lambda]^\mu$ for some $a \in \mathscr{P}$ we have $a \subseteq A\}$. If $\mu = \lambda$ we may omit it. Let ⫯_λ mean $\text{⫯}_{\lambda,\aleph_1}$ and $\text{⫯} = \text{⫯}_{\aleph_1}$.

4.5 Claim. *1)* $⫯_{\lambda,\mu,\theta} = \ dcf_{\langle\{\emptyset\},[\lambda]^{<\theta},[\lambda]^{<\mu}\rangle}(\lambda)$ *when* $\theta \le \mu \le \lambda$.
2) $ecf_{\mu,\theta,\sigma}(\lambda,\theta) = \min\{|\mathscr{P}| : \mathscr{P} \subseteq [\lambda]^\theta$ *and for every* $A \in [\lambda]^\mu$ *for some* $a \in \mathscr{P}$ *we have* $|A \cap a| \ge \sigma\}$.
Now we can phrase the analog of 4.1 for dcf.

4.6 Lemma. *Assume* $\theta < \mu < \lambda \le \lambda^*, J$ *an ideal on* θ *and assume*

$\otimes^J_{\theta,\mu,\lambda,\lambda^*}$ *if* $n < \omega, \mathfrak{a}_i \in [Rang\ \cap \lambda^+ \backslash \mu^+]^n$ *for* $i < \theta$ *then* $\{a \in J : max\ pcf \bigcup\limits_{i \in a} \mathfrak{a}_i \le \lambda^*\}$ *is generated by* $\le \mu$ *sets.*[2]

Then there is a set H *such that*

- (a) H *a st of partial functions from* θ *to* $[\lambda]^{\le\mu}$
- (b) $|H| \le \lambda^*$
- (c) *for every function* $g : \theta \to \lambda$ *we can find* h *and* $\bar{\mathfrak{a}} = \langle \mathfrak{a}_i : i < \theta\rangle$ *such that*
  - (i) $\mathfrak{a}_i$ *is a finite set of regular cardinals from* $(\mu, \lambda]$
  - (ii) h *is a function from* θ *to* $[\lambda]^{\le\mu}$ *such that* $\bigwedge\limits_i g(i) \in h(i)$
  - (iii) *for any* $n < \omega$ *and* $a \subseteq \theta$: *if* $(\forall i \in a)[|\mathfrak{a}_i| \le n]$ *and max pcf*$(\bigcup\limits_{i\in a} \mathfrak{a}_i) \le \lambda^*$ *then for some* $b, a \subseteq b \subseteq \theta$ *we have* $h \restriction b \in H$.

4.7 Claim. *Assume:*

$(*)_0$ $\aleph_0 < \aleph_{\alpha(*)} \le ⫯$ *and:* $\aleph_{\alpha(*)} < \aleph_{\omega_2}$ *or at least*

$$cov(\aleph_{\alpha(*)}, \aleph_2, \aleph_2, \aleph_1) \le ⫯$$

$(*)_1$ $\mathfrak{a} \subseteq\ Reg \cap ⫯ \backslash \aleph_{\alpha(*)+1}\ \ \&\ \ |\mathfrak{a}| \le \aleph_0 \Rightarrow |pcf(\mathfrak{a})| \le \aleph_{\alpha(*)}$

$(*)_2$ *if* $\lambda_i \in (\aleph_1, ⫯) \cap\ Reg$ *for* $i < \omega_1$ *then for some* $a \in [\omega_1]^{\aleph_1}$, *for every* $b \in [a]^{\aleph_0}$ *we have max pcf*$(\{\lambda_i : i \in b\}) \le ⫯$ *then* $⫯_{⫯} = ⫯$.

4.8 Remark. 1) This means that the conclusion holds except when some dubious statements on pcf, ones which have high consistency strength (or are inconsistent) and ⫯ is somewhat large.
2) There are obvious monotonicity properties and $\text{ecf}_{\bar{J}}(\lambda, < \mu) \le\ \text{dcf}_{\bar{J}}(\lambda, < \mu)$.

Proof. Let $\theta = \aleph_1, \mu = \aleph_{\alpha(*)}, \lambda = ⫯, \lambda^* = ⫯, J = [\omega_1]^{\le\aleph_0}$. Apply 4.2, more exactly the variant 4.2(2). The assumption $\otimes^J_{\theta,\mu,\lambda,\lambda} = \otimes^J_{\aleph_1,\aleph_{\alpha(*)},⫯,⫯}$ holds by $(*)_1$. So let $H \subseteq \{h : h : \aleph_1 \to [⫯]^\mu\}, |H| = \lambda^* = \lambda$ be as in the conclusion there. Let

---

[2] see 4.2(3)

$\chi = \beth_7^+, \mathfrak{B} \prec (\mathscr{H}(\chi), \in, <^*_\chi), |\mathfrak{B}| = \lambda, \lambda + 1 \subseteq \mathfrak{B}, H \in \mathfrak{B}$. We want to show $\mathscr{P} = \mathfrak{B} \cap [\lambda]^{\aleph_0}$ exemplifies $\text{⫯}_\lambda = \lambda$. So let $g : \theta \to \lambda$, so there are $h, \langle\langle\lambda^n_i : n < n_i\rangle : i < \omega_1\rangle$, as there. Let n^* be such that $B_0 =: \{i < \omega_1 : n_i = n^*\}$ is uncountable. By using $(*)_2$ n times we can find an uncountable $B \subseteq B_0 (\subseteq \omega_1)$ such that

$$(*) \quad a \subseteq B \ \ \& \ \ a \in J \Rightarrow \ \max \mathrm{pcf}\{\lambda^n_i : n < n^*, i \in a\} \leq \lambda.$$

So for every $a \in [B]^{\aleph_0}$, for some $b \in [\omega_1]^{\aleph_0}$ we have $a \subseteq b$ and $h \restriction b \in H \subseteq \mathfrak{B}$.

Let for a set $b \in H(\chi)$ of ordinals, f_b be the $<^*_\chi$-first one- to-one function from $|b|$ onto b, let $g'(i) = f^{-1}_{h(i)}(g(i))$, so g' is a function from $\theta = \aleph_1$ to $\mu = \aleph_{\alpha(*)}$ (as $|h(i)| \leq \mu$). Now $\mathrm{cov}(\aleph_{\alpha(*)}, \aleph_2, \aleph_2, \aleph_1) \leq \text{⫯}$ so $\mathrm{cov}(\aleph_{\alpha(*)}, \aleph_2, \aleph_2, \aleph_1) \leq \lambda$ (the only property of $\alpha(*)$ we use) so there is $\mathscr{P}' \subseteq [\aleph_{\alpha(*)}]^{\leq \aleph_1}, |\mathscr{P}'| \leq \lambda$ exemplifying this and without loss of generality $\mathscr{P}' \subseteq \mathfrak{B} \ \ \& \ \ \mathscr{P}' \in \mathfrak{B}$. So the set $\{g'(i) : i \in B\}$ is included in a countable union of members of $\mathscr{P}'$, so for some $Y \in \mathscr{P}'$ (so $Y \in [\aleph_{\alpha(*)}]^{\leq \aleph_1}, Y \in \mathfrak{B}$) we have $B^* =: \{i \in B : g'(i) \in Y\}$ is uncountable.

Define h':

$$\mathrm{Dom}(h') = \omega_1, h'(i) = \{\alpha \in h(i) : f_{h(i)}(\alpha) \in Y\}.$$

So h' is a function from ω_1 to $[\lambda]^{\leq \aleph_1}$ (as $|Y| \leq \aleph_1$) and $i \in B^* \Rightarrow g(i) \in h'(i)$; remember $a \in [B^*]^{\leq \aleph_0} \Rightarrow (\exists b)[a \subseteq b \subseteq \omega_1 \ \ \& \ \ h \restriction b \in \mathfrak{B}]$.

Let $Z =: \{(i, f^{-1}_{h'(i)}(g(i))) : i \in B\}$, it is a subset of $\omega_1 \times \omega_1$ of cardinality λ, but $\text{⫯} = \lambda, \lambda + 1 \subseteq \mathfrak{B}$, so for some infinite $z \in \mathfrak{B}, z \subseteq Z$. Let $z_0 = \{i < \omega_1 : \bigvee\limits_j (i, j) \in z\}$, so $z_0 \in \mathfrak{B}, z_0 \in [\omega_1]^{\aleph_0}$ and even $z_0 \subseteq B^*$, hence $h' \restriction z_0 \in \mathfrak{B}$. So as $h' \restriction z_0 \in \mathfrak{B}$ and $\{(i, f^{-1}_{h'(i)}(g(i))) : i \in z_0\} \in \mathfrak{B}$ also $g \restriction z_0 \in \mathfrak{B}$, so $\mathrm{Rang}(g \restriction z_0) \in B$, so $\mathrm{Rang}(g \restriction z_0) \in \mathscr{P}$ and we are done. $\square_{4.7}$

4.9 Definition.

$$\begin{aligned} St^3_{\lambda,\kappa} = \min\big\{|\mathscr{P}| : &(a) \quad \mathscr{P} \subseteq [\lambda]^{\aleph_0} \\ &(b) \quad (\forall A \in [\lambda]^\kappa)(\exists b \in \mathscr{P})(b \cap A \text{ infinite}) \\ &(c) \quad \mathscr{P} \text{ is } AD \text{ which means } a \neq b \in \mathscr{P} \Rightarrow a \cap b \text{ finite}\big\} \end{aligned}$$

(main case $\kappa = \aleph_1$).

4.10 Definition.

$$\begin{aligned} St^4_{\lambda,\kappa} = \min\big\{|\mathscr{P}| : &(a) \quad \mathscr{P} \subseteq [\lambda]^\omega \\ &(b) \quad (\forall A \in [\lambda]^\kappa)(\exists b \in \mathscr{P})(b \cap A \text{ infinite}) \\ &(c) \quad \sup\{\mathrm{otp}(a) : a \in \mathscr{P}\} < \omega_1\big\}. \end{aligned}$$

4.11 Definition.

$$
\begin{aligned}
St^5_{\lambda,\kappa} = \min\big\{\mathscr{P} :\, &(a) \quad \mathscr{P} \subseteq [\lambda]^\omega \\
&(b) \quad (\forall A \in [\lambda]^\kappa)(\exists b \in \mathscr{P})(b \cap A \text{ infinite}) \\
&(c) \quad \text{the } BA \text{ of subsets of } \lambda \text{ which } \mathscr{P} \text{ and the singletons} \\
&\qquad \text{generate is superatomic of rank } < \omega_1\big\}.
\end{aligned}
$$

4.12 Fact: $\mathrm{dcf}_{\kappa,\aleph_0,\aleph_0} \le St^\ell_{\lambda,\kappa}$.

* * *

4.13 Claim. *1) Given $\lambda \ge \kappa = cf(\kappa) > \aleph_0$, the following cardinals are equivalent for $k < \omega, k > 0$:*

(a) $ecf_{\kappa,\aleph_0,\aleph_0}(\lambda)$

$(b)_k$ $min\{|\mathscr{F}|$:

- (i) *$\mathscr{F}$ is a family of partial functions f from λ to k*
- (ii) $f \in \mathscr{F} \Rightarrow |Dom\ f| = \aleph_0$,
- (iii) *$f \in \mathscr{F}, \ell < k \Rightarrow f^{-1}(\{\ell\})$ is infinite*
- (iv) *if $\langle A_0, \ldots, A_{k-1}\rangle$ are pairwise disjoint subsets of λ each of cardinality κ then for some $f \in \mathscr{F}$ we have $\ell < k \Rightarrow f^{-1}(\{\ell\}) \cap A_\ell$ is infinite$\}$*

$(c)_k$ *like $(b)_k$ replacing (iii), (iv) by*

- $(iii)^+$ *if $\langle \alpha_{\varepsilon,\ell} : \varepsilon < \kappa, \ell < k\rangle$ is a sequence of ordinals, with no repetitions then for infinitely many $\varepsilon < \kappa$, for some $\ell < k, f(\alpha_{\varepsilon,\ell}) = \ell$ (so $\alpha_{\varepsilon,\ell} \in$ Dom(f)).*

Proof. Let $\lambda^b_\kappa, \lambda^c_k$ be the cardinal from $(b)_k, (c)_k$ respectively and λ^* the cardinal from (a). Clearly $\lambda^b_k \le \lambda^b_{k+1}, \lambda^c_k \le \lambda^c_{k+1}, \lambda^b_k \le \lambda^c_k, \lambda^c_1 = \lambda^b_1 = \lambda^*$. So it suffices to prove $\lambda^c_k \le \lambda^*$, assume $\mathscr{P}$ exemplifies $\lambda^* = \mathrm{ecf}_{\kappa,\aleph_0,\aleph_0}(\lambda)$ by 4.5 (2). If $\lambda^* \ge 2^{\aleph_0}$ let $\mathscr{F}^* = \{f$: for some $a \in \mathscr{P}, f$ is a function from a to k such that $\ell < k \Rightarrow |f^{-1}(\{\ell\})| = \aleph_0\}$ clearly it exemplifies $\lambda^c_k \le |\mathscr{F}^*| = 2^{\aleph_0} \times \lambda^* = \lambda^*$. So assume $\lambda^* < 2^{\aleph_0}$, hence $\lambda^* < 2^{\aleph_0}$ and let $\bar\eta = \langle \eta_i : i < \lambda^*\rangle$ be a sequence of pairwise distinct members of ${}^\omega 2$. Let $\bar g = \langle g_\ell : \ell < k\rangle$ be such that: $g_\ell : \lambda \to \lambda$ and $(\forall \alpha_0 \ldots \alpha_{k-1} < \lambda)(\exists \beta < \lambda)[\bigwedge\limits_{\ell<k} g_\ell(\beta) = \alpha_\ell]$. Let $\mathscr{P}' = \{a^{\bar g} : a \in \mathscr{P}\}$ where $a^{\bar g} = \{g_\ell(x) : \ell < k, x \in a\}$ and let $\mathscr{F} = \{f_{b,h} : b \in \mathscr{P}$ and for some n we have $h \in {}^{n}2)k$ and $\ell < k \Rightarrow |f^{-1}_{b,h}(\{\ell\}) = \aleph_0\}$ where $f_{b,h}$ is the function with domain $b^{\bar g}, f_{b,h}(i) = h(\eta_i \restriction n)$ where $h \in {}^{({}^{n}2)}k$.

Clearly $\mathscr{F}$ has the right cardinality and form. Let us show that it satisfies the main requirement: let $\langle A_0, \ldots, A_{k-1}\rangle$ be a sequence of subsets of λ each of cardinality κ. Let $A_\ell = \{\gamma_{\ell,\varepsilon} : \varepsilon < \kappa\}$ (no repetition). Let $\gamma_\varepsilon < \lambda$ be such that

$\bigwedge\limits_{\ell<k} g_\ell(\gamma_\varepsilon) = \gamma_{\ell,\varepsilon}$. For each ε for some $n(\varepsilon)$ we have: $\langle \eta_{\gamma_{\ell,\varepsilon}} \restriction n(\varepsilon) : \ell < k\rangle$ is with no repetitions. As $\kappa = \mathrm{cf}(\kappa) > \aleph_0$ without loss of generality for some $\bar\nu = \langle \nu_\ell : \ell < k\rangle$ and $n(*)$ we have $\varepsilon < \kappa \Rightarrow \eta_{\gamma_{\ell,\varepsilon}} \restriction n(*) = \nu_\ell, \varepsilon < \kappa \Rightarrow n(\varepsilon) = n(*)$. Now by the choice of $\mathscr{P}$ for some $a \in \mathscr{P}, W = \{\varepsilon : \gamma_\varepsilon \in a\}$ is infinite. Let $b = a_{\bar g}$, let $h : {}^{n(*)}2 \rightarrow k$ be such that $h(\nu_\ell) = \ell$, now $f_{b,h} \in \mathscr{F}$ is as required. $\Box_{4.13}$

4.14 Claim. *Assume $\lambda_{n+1} \geq \mathrm{ecf}_{\kappa,\aleph_0,\aleph_0}(\lambda_n, \aleph_0)$ for $n < \omega, \lambda_0 \geq \kappa = \mathrm{cf}(\kappa) > \aleph_0$. Then there is a Boolean Algebra B of cardinality $\sum\limits_{n<\omega} \lambda_n$ into which the free Boolean algebra generated by $\sum\limits_{n<\omega} \lambda_n$ elements can be embedded but such that there is no homomorphism from B onto the free Boolean Algebra generated by κ elements.*

4.15 Remark. 1) So we can find quite many pairs $\lambda = \sum\limits_{n<\omega} \lambda_n$ and κ as required in 4.14 (or 4.15). E.g. any $\lambda = \lambda_n > \beth_\omega$ and large enough $\kappa = \mathrm{cf}(\kappa) \leq \beth_\omega$ is as required by [Sh 460]; also $\lambda = \lambda_n = \aleph_{\kappa^{+4}} > \kappa = \mathrm{cf}(\kappa) > \aleph_0$ is as required by [Sh 400, §4].
2) On the problem see Fuchino, Shelah, Soukup [FShS 543], [FShS 544].
3) From the proof we can strengthen the last phrase in the conclusion to "no homomorphism from B into $Fr(\kappa)$ with range of cardinality κ". Similarly in 4.17.

Proof. Let $\mathscr{F}_\alpha = \{f^n_\alpha : \alpha < \lambda_{n+1}\}$ be as required in clause $(b)_2$ of 4.13 (exists as $\lambda_{n+1} \geq \mathrm{ecf}_{\kappa,\aleph_0,\aleph_0}(\lambda_n)$).

Remember the variety of Boolean rings has the operation $x \cup y, x \cap y, x - y$ and constant 0 (but no 1 and no $-x$), so any ideal of a Boolean algebra is a Boolean ring and if the ideal is maximal, the Boolean algebra is definable in the Boolean ring.

Let B_0 be the Boolean ring freely generated by $\{x^0_i : i < \lambda_0\}$. Let B_1 be the Boolean ring generated by $B_0 \cup \{x^1_i : i < \lambda_1\}$ freely except:

(a) the equations which holds in B_0
(b) $x^1_{2\alpha} \cap x^0_i = 0$ if $f^n_\alpha(i) = 0$ (so $i \in \mathrm{Dom}(f^n_\alpha)$)
(c) $x^1_{2\alpha} \cap x^0_i = x^0_i$ if $f^n_\alpha(i) = 1$ (so $i \in \mathrm{Dom}(f^n_\alpha)$).

(Why $x^1_{2\alpha}$ and not x^1_α? to embed $Fr(\sum\limits_{n<\omega} \lambda_n)$ into the Boolean Algebra we are constructing.)

Similarly let B_{n+1} be the Boolean ring generated by $B_n \cup \{x^{n+1}_\alpha : \alpha < \lambda_{n+1}\}$ freely except

(a) the equations which holds in B_n
(b) $x^{n+1}_{2\alpha} \cap x^n_i = 0$ if $f^n_\alpha(i) = 0$ (if $i \in \mathrm{Dom}(f^n_\alpha)$)
(c) $x^{n+1}_\alpha \cap x^n_i = x^n_i$ if $f^n_\alpha(i) = 1$ (if $i \in \mathrm{Dom}(f^n_\alpha)$).

Now $B_\omega = \bigcup\limits_{n<\omega} B_n$ is a Boolean ring and let B be the Boolean algebra for which B_ω is a maximal ideal. Assume f is a homomorphism from B onto $Fr(\kappa)$, the

Boolean algebra freely generated say by $\{z_i : i < \kappa\}$. Now B is generated by $\{x^n_\alpha : n < \omega, \alpha < \lambda_n\}$. So as f is onto, for some n, for every $\zeta < \kappa$ for some $\alpha, f(x^n_{\alpha,\sigma_0}) \notin \langle z_\varepsilon : \varepsilon < \zeta \rangle_{Fr(\kappa)}$. By the Δ-system lemma, we can find a stationary $S \subseteq \kappa$, Boolean term $\sigma_1 = \sigma_1(x_0, \dots, x_{n(*)-1}), m(*) < n(*)$ ordinals $\varepsilon(0) < \dots < \varepsilon(m(*)-1) < \min(S)$, and for each $\zeta \in S$, ordinals $\varepsilon(m(*),\zeta) < \dots < \varepsilon(n(*)-1,\zeta)$ all in the interval $[\zeta, \min(S \backslash (\zeta+1))]$ and $\alpha_\zeta < \lambda_n$ such that $f(x^n_{\alpha_\zeta,\sigma}) = \sigma_1(z_{\varepsilon(0)}, \dots, z_{\varepsilon(m(*)-1)}, z_{\varepsilon(m(*),\zeta)}, \dots, z_{\varepsilon(n(*)-1,\zeta)})$, where all the $n(*)$ variables are needed in the term σ_1.

Let $S = \{\zeta(i) : i < \kappa\}$ with $\zeta(i)$ increasing in i, let f be the function $f(\alpha_{\zeta(2i+1)}) = 1, f(\alpha_{\zeta(2i)}) = 0$. So for some $\alpha < \lambda_{n+1}$ for $\ell = 0, 1$ the sets $W_\ell =: \{\alpha_{\zeta(2i+\ell)} : f^n_\alpha(\alpha_{\zeta(2i+\ell)}) = \ell\}$ are infinite. So $z^* = f(x^{n+1}_{2\alpha}) \in Fr(\kappa)$ satisfies

$(*)_0$ $\alpha_\zeta \in W_0 \Rightarrow Fr(\kappa) \models z^* \cap f(x^n_{\alpha_\zeta}) = 0$

$(*)_1$ $\alpha_\zeta \in W_1 \Rightarrow Fr(\kappa) \models z^* \cap f(x^n_{\alpha_\zeta}) = f(x^n_{\alpha_\zeta})$.

But for some finite $u \subseteq \kappa, z^* \in \langle z_\gamma : \gamma \in u \rangle Fr(\kappa)$, so there is $\alpha_{\zeta(2i_0)} \in W_0$, such that u is disjiont to $\{\varepsilon(m(*), \zeta(2i_0)), \dots, \varepsilon(n(*)-1, \zeta(2i_0+1))\}$ and there is $\alpha_{\zeta(2i_1+1)} \in W_1$ such that u is disjoint to $\{\varepsilon(m(*), \zeta(2i_1+1)), \dots, \varepsilon(n(*)-1, \zeta(2i_1+1))\}$. For them $(*)_0, (*)_1$ gives a contradiction. So B is a Boolean algebra, of cardinality $\leq \lambda$ (as it is generated by $\{x^n_\alpha : \alpha < \lambda_n, n < \omega\}$), we can embed into it the free Boolean algebra with λ generators $\{x^n_{2\alpha+1} : \alpha < \lambda_n, n < \omega\}$) with no homomorphism onto the free Boolean algebra with κ generators. $\square_{4.14}$

4.16 Definition. Let B^{fcf}_μ be the Boolean Algebra of finite and cofinite subsets of μ.

4.17 Claim. *Assume* $\lambda_{n+1} \geq \text{ecf}_{\kappa,\aleph_0,\aleph_0}(\lambda_n, \aleph_0), \kappa = \text{cf}(\kappa) > \aleph_0$ *and* $\lambda = \sum_n \lambda_n$. *Then there is a Boolean Algebra* B *of cardinality* λ *into which* B^{fcf}_λ *can be embedded but such that there is no homomorphism from* B *onto* B^{fcf}_κ.

Proof. Let

$$\mathscr{P}_n = \{a^n_\alpha : \alpha < \lambda_{n+1}\} \subseteq [\lambda_n]^{\aleph_0}$$

exemplifies $\lambda_{n+1} \geq \text{ecf}_{\kappa,\aleph_0,\aleph_0}(\lambda_n, \aleph_0)$ by 4.5(2). We define by induction on n a countable subset x^n_α of $n \times \lambda$ for each $\alpha < \lambda_n$. For $n = 0$ let $x^0_\alpha = \{(0, \alpha)\}$. For $n+1$ let

$$x^{n+1}_{2\alpha} = \{(n+1, 2\alpha)\} \cup \bigcup_{\beta \in a^n_\alpha} x^n_\beta \text{ and } x^{2n}_{2\alpha+1} = \{(n+1, 2\alpha+1)\}.$$

Let B be the Boolean Algebra of subsets of $\omega \times \lambda$ generated by $\{x^n_\alpha : \alpha < \lambda_n, n < \omega\}$.

Clearly $|B| \leq \sum_n \lambda_n = \lambda$, also $\{x^n_{2\alpha+1} : \alpha < \lambda, n < \omega\}$ generate a subalgebra isomorphic to B^{fcf}_λ hence $|B| \geq \lambda$ (so $|B| = \lambda$) and B^{fcf}_λ can be embedded into B.

Lastly, suppose g is a homomorphism from B onto B^{fcf}_κ. Let $z_\zeta =: \{\zeta\} \in B^{fcf}_\kappa$. For each $\zeta < \kappa$ for some $n(\zeta) < \omega, \alpha(\zeta) < \lambda_n$ we have $g(x^{n(\zeta)}_{\alpha(\zeta)}) \notin \langle z_\zeta : \varepsilon < \zeta\rangle_{B^{fcf}_\kappa}$, so for some stationary $S \subseteq \kappa, [\zeta \in S \Rightarrow n(\zeta) = n(*)]$. If $g(x^n_\alpha) \in B^{fcf}_\kappa$ is infinite then $\{g(x) : x \in B \text{ and } g(x) \le g(x^n_\alpha)\} = \{g(x \cap x^n_\alpha) : x \in B\}$ is countable so g is not onto B, a contradiction. So possibly shrinking S without loss of generality $\langle g(x^{n(*)}_{\alpha(\zeta)}) : \zeta \in S\rangle$ is a Δ-system of finite subsets of κ with heart called w.

For some $\beta < \lambda_{n+1}$ the set $u = \{\zeta \in S : \alpha(\zeta) \in a^{n+1}_\beta\}$ is infinite, clearly $\zeta \in u \Rightarrow x^n_{\alpha(\zeta)} \le x^{n+1}_{2\beta}$ hence $g(x^{n(*)}_{\alpha(\zeta)}) \le g(a^{n+1}_{2\beta})$ hence $g(x^{n+1}_{2\beta})$ is infinite hence it is co-finite, contradicting an earier statement. $\square_{4.17}$

4.18 Definition.

$$
\begin{aligned}
St^6_{\lambda,\kappa} = \min\big\{\lambda^* :& \text{there is } \mathscr{P} \subseteq [\lambda]^{\aleph_0} \text{ such that} \\
&(i) \quad \mathscr{P} \text{ is } \aleph_2\text{-free i.e. if } a_i \in P \text{ for } i < i^* < \aleph_2 \text{ are disjoint} \\
&\qquad \text{then for some finite } b_i \subseteq a_i, \langle a_i \backslash b_i : i < i^*\rangle \\
&\qquad \text{are pairwise disjoint} \\
&(ii) \quad \text{for every } f : \kappa \to \lambda \text{ for some } a \in P \\
&\qquad\qquad (\exists^\infty \alpha < \kappa)(f(\alpha) \in a)\big\}.
\end{aligned}
$$

4.19 Claim. *Assume* $\lambda_{n+1} \le St^6_{\lambda_n,\kappa}$ *for* $n < \omega, \lambda = \sum\limits_{n<\omega} \lambda_n$. *Then there is a Boolean Algebra* B *as in the previous claim which is superatomic.*

Proof. Like the previous claim.

§5 More on independence

5.1 Claim. *If* $|\mathrm{pcf}(\mathfrak{a})| \geq \mu = \mathrm{cf}(\mu) > \lambda^+ > |\mathfrak{a}|^+, \sigma < \lambda$ *and* $\mathrm{cf}(\chi) = \mu, \chi = |\chi \cap \mathrm{pcf}(\mathfrak{a}|, \chi$ *is* J^{bd}_λ*-inaccessible,* *then for any* $\chi_0 < \chi$ *there is* $\mathfrak{b} \subseteq \mathrm{pcf}\, \sigma$*-complete* $(\mathfrak{a}) \cap (\chi_0, \chi) \backslash \lambda, |\mathfrak{b}| < \lambda$*, such that* $pcf_{\sigma\text{-complete}}(\mathfrak{b}) \cap \chi$ *is unbounded in* χ*.*

Proof. We use [Sh 430], 6.7F(5) (localization for $\mathrm{pcf}_{\sigma\text{-complete}}$), [Sh 420],§1. $\square_{5.1}$

5.2 Claim. *Assume* $IND(\langle J^{bd}_{\lambda_\varepsilon} : \varepsilon < \varepsilon(*)\rangle)$ *and* $\lambda = \sum\limits_{\varepsilon < \varepsilon(*)} \lambda_\varepsilon$*. If* $\mu > \lambda$ *and* $\theta_i \in Reg \cap \mu \backslash \lambda$ *for* $i < \lambda$*, then for some* ε *we can find* $\mathfrak{c} \subseteq \mu \cap \mathrm{pcf}\{\theta_i : i < \lambda\}, |\mathfrak{c}| \leq \lambda$ *and* $\mathfrak{b}_\tau \in J_{\leq \tau}[\{\theta_i : i < \lambda\}]$ *for* $\tau \in \mathfrak{c}$ *such that*

$(*)$ *there is no* $a \in [\lambda]^{\lambda_\varepsilon}$ *such that* $[\tau \in \mathfrak{c} \Rightarrow (\forall^{|\mathfrak{a}|} i \in a)\theta_i \notin \mathfrak{b}_\tau]$
$((\forall^{|\mathfrak{a}|} i \in a)$ *means: for all but* $< |a|$ *members* i *of* a*).*

Proof. Repeat the proof of **?**.
−> scite{3.x} undefined

5.3 Remark. If we deal with a normal ideal it seems that, for a given $\chi \in \mathrm{acc}(c\ell(\mathrm{pcf}(\mathfrak{a})))$, we can get long intervals of λ with χ being J^{bd}_λ-inaccessible.

5.4 Claim. *1) If* $IND(\lambda, \kappa)$*, then for every* $\mathfrak{a} \subseteq Reg$*,* $\lambda \leq |\mathfrak{a}| < \min(\mathfrak{a})$ *for some* $\mathfrak{b} \subseteq \mathfrak{a}, \kappa < |\mathfrak{b}| \leq \lambda, \Pi(\mathfrak{b})/[\mathfrak{b}]^{\leq \kappa}$ *has true cofinality (so if* λ *is minimal for this* κ*, this holds for any* $\kappa' < \lambda$*).*
2) If $IND(\langle J^{bd}_{\lambda_n} : n < \omega\rangle), \lambda_n < \lambda_{n+1}, \lambda_n$ *regular,* $|\mathfrak{a}| < \lambda_0, |\mathfrak{a}| < \min(\mathfrak{a})$ *and*

$$\mu = \sup_{n<\omega} \min\{|\mathscr{P}| : \mathscr{P} \subseteq [\lambda_n]^{|\mathfrak{a}|^+} \textit{ and } (\forall A \in [\lambda_n]^{\lambda_n})(\exists B \in \mathscr{P})[B \subseteq A]\}$$

then $|pcf(\mathfrak{a}) \leq \mu$*.*
3) Assume $\sigma < \theta < \lambda_n\, (n < \omega)$*. (We can guess filters: which are* $(< \theta)$*-based.)* $IND(\langle J_n : n < \omega\rangle), J_n$ *an ideal on* λ_n *and* μ *satisfies:* $\bigwedge\limits_n \mu > \kappa_n$ *and*

$(*)_{\mu, J_n}$ *there is a set* $\mathscr{E}, |\mathscr{D}| \leq \mu$*, each member of* $\mathscr{E}$ *is an ideal on some bounded subset of* κ_n *such that:*

- $\otimes$ *if* $Y \in J^+_n$ *(so* $Y \subseteq \lambda_n$*), and* I *is a* $(< \theta)$*-based* σ*-complete ideal on* Y *generated by* $\leq \mu$ *sets then for some* $I' \in \mathscr{E}$*, we have* $(Dom\ I') \cap Y \in I^+$ *and* $Dom(I') \backslash Y \in I', I' \restriction (Y \cap\ Dom(I')) \supseteq I \restriction (Y \cap\ Dom(I'))$*.*

Then for $\mathfrak{a} \subseteq Reg$*,* $(pcf_{\sigma\text{-complete}}(\mathfrak{a})) \leq \mu$*.*

Proof. 1) Straight.
2) Suppose not. Let $\langle \tau_{i+1} : i < \mu^+\rangle$ be the first μ^+ members of $\mathrm{pcf}(\mathfrak{a})$ listed in

increasing order. Let $\tau_\delta = \bigcup\limits_{i<\delta} \tau_{1+i}$ for limit $\delta \le \mu^+$. For each limit $\delta < \mu^+$ for some $n = n_\delta < \omega, \tau_\delta$ is $\{J^{bd}_{\lambda_n}\}$-inaccessible (by 3.13?). So for some $n(*) < \omega, \{\delta < \mu^+ : n_\delta = n(*)\}$ is stationary, hence

(∗) for no $\theta_\alpha \in$ Reg $\cap\, \tau_{\mu^+}$ for $\alpha < \lambda_{n(*)}$, do we have

$$\prod_{\alpha<\lambda_{n(*)}} \theta_\alpha / J^{bd}_{\lambda_{n(*)}} \text{ is } \tau_{\mu^+}\text{-directed.}$$

By [Sh 420, §1] we can find $\langle C_\alpha : \alpha \in S\rangle, S \subseteq \mu^+, C_\alpha \subseteq \alpha$, $\mathrm{otp}(C_\alpha) \le \lambda_{n(*)}, [\beta \in C_\alpha \Rightarrow \beta \in S \ \ \& \ \ C_\beta = C_\alpha \cap \beta]$ and $\mathrm{otp}(C_\alpha) = \lambda_{n(*)} \Rightarrow \alpha = \sup(C_\alpha)$ and $\{\alpha \in S : \mathrm{otp}(C_\alpha) = \lambda_{n(*)}\}$ is stationary. Now we imitate [Sh 400, §2].
3) Similar to 2). $\square_{5.4}$

§6 Odds and ends

As in [Sh 430, §6] this section is dedicated to things I forgot to say. We repeat and elaborate older things [Sh 430], **?**, **?**, **?**, [Sh 410], 3.7.

$\Rightarrow$ scite{6.6D} undefined
$\Rightarrow$ scite{6.6E} undefined
$\Rightarrow$ scite{6.6F} undefined

6.1 Claim. *Suppose D is a σ-complete filter on $\theta =$ cf(θ) such that $[\alpha < \theta \Rightarrow \theta \backslash \alpha \in D], \sigma$ is regular $> \kappa^+ + |\alpha|^\kappa$ for $\alpha < \sigma$, and for each $\alpha < \theta, \bar{\beta} = \langle \beta^\alpha_\epsilon : \epsilon < \kappa \rangle$ is a sequence of ordinals. Then for every $X \subseteq \theta, X \neq \emptyset$ mod D there is $\langle \beta^*_\epsilon : \epsilon < \kappa \rangle$ (a sequence of ordinals) and $w \subseteq \kappa$ such that:*

- *(a) $\epsilon \in \kappa \backslash w \Rightarrow \sigma \leq$ cf(β^*_ϵ) $\leq \theta$,*
- *(b) $B =: \{\alpha \in X :$ if $\epsilon \in w$ then $B^\alpha_\epsilon = \beta^*_\epsilon$ and: if $\epsilon \in \kappa \backslash w$ then β^α_ϵ is $< \beta^*_\epsilon$ but $> \sup\{\beta^*_\zeta : \zeta < \kappa, \beta^*_\zeta < \beta^*_\epsilon\}\}$ is $\neq \emptyset$ mod D*
- *(c) if $\beta'_\epsilon < \beta^*_\epsilon$ for $\epsilon \in \kappa \backslash w$ then $\{\alpha \in B :$ if $\epsilon \in \kappa \backslash w$ then $B'_\epsilon < \beta^\alpha_\epsilon\} \neq \emptyset$ mod D.*

6.2 Remark. 1) Of course, we can replace κ by any set of this cardinality.
2) May look at [Sh 620], §7, there more is said concerning 6.1.

Proof. Let $f_\alpha : \kappa \rightarrow$ Ord be $f_\alpha(i) = \beta^\alpha_i$.

Let χ be large enough. We choose by induction on $i < \sigma$, a model N_i such that:

$N_i \prec (\mathscr{H}(\chi), \in, <^*_\chi)$;
$\|N_i\| \leq 2^\kappa + |i|^\kappa$;
$2^\kappa \subseteq N_0$;
$\kappa, \sigma, \theta \in N_0, \langle f_\alpha : \alpha < \theta \rangle \in N_0$;
$i < j \Rightarrow N_i \prec N_j$;
$\langle N_j : j \leq i \rangle \in N_{i+1}$;
N_i increasing continuous.

Let $\delta_i =: \min(\cap \{B : B \in N_i \cap D\})$, now δ_i is well defined (as D is σ-complete and $\sigma > 2^\kappa + |i|^\kappa \geq \|N_i\|$, hence the intersection is in D). Now $\delta_i \geq \sup(\theta \cap N_i)$. As $\alpha \in \theta \cap N_i \Rightarrow \theta \backslash \alpha \in D \cap N_i$ and as $\delta_i \in N_{i+1}$ (as $\{N_i, D, \theta\} \in N_{i+1}$) clearly $\langle \delta_i : i < \sigma \rangle$ is strictly increasing continuous. We define for $i < \sigma$, a function $g_i \in {}^\kappa \chi$ by

$$g_i(\zeta) = \min(N_i \cap \chi \backslash f_{\delta_i}(\zeta))$$

(it is well defined as $\bigcup\limits_{\alpha < \theta} (f_\alpha(i) + 1) \in N_0 \prec N_i$). Clearly $E =: \{\alpha < \sigma : N_\alpha \cap \sigma = \alpha\}$ is a club of σ, and as $(\forall \alpha < \sigma)[|\alpha|^\kappa < \sigma]$ clearly $\alpha < \beta \in E \ \& \ a \subseteq N_\alpha \ \& \ |a| \leq \kappa \Rightarrow a \in N_\beta$. Now $i \in E$, cf$(i) = \kappa^+$ implies $N_i = \bigcup\limits_{j<i} N_j$ and Rang$(g_i) \subseteq \bigcup\limits_{j<i} N_j$ hence $\bigvee\limits_{j<i} [\text{Rang}(g_i) \subseteq N_j]$; but by the previous sentence every subset of N_j of cardinality $\leq \kappa$ belongs to N_i, hence $g_i \in \bigcup\limits_{j<i} N_j$. So by Fodor Lemma for some

stationary subset S of $\{i \in E : \mathrm{cf}(i) = \kappa^+\}$ and some $g^* : \kappa \rightarrow \sigma$ and some $u \subseteq \kappa$ and some $i(*) < \sigma$ we have: $[i \in S \Rightarrow g_i = g^*], (\forall i \in S)(\forall \zeta < \kappa)[f_{\delta_i}(\zeta) = g^*(\zeta) \Leftrightarrow \zeta \notin u]$ and $g^* \in N_{i(*)}$; note $u \in N_0 \subseteq N_{i(*)}$ as $u \subseteq \kappa$ and we can assume $i(*) < \min(S)$ and $i(*) \in E$.

Let $w =: \kappa \backslash u, \beta^*_i =: g^*(i)$ for $i < \kappa$ now we show that $w, \langle \beta^*_i : i < \kappa \rangle$ are as required.

Clause (b):

The set B is defined from: $\langle B^*_i : i < \kappa \rangle$ and w and $\bar{f} = \langle f_\alpha : \alpha < \theta \rangle$. As all of them belong to $N_{i(*)}$ clearly $B \in N_{i(*)}$, so if $B = \emptyset \mod D$ then $(\theta \backslash B) \in D \cap N_{i(*)}$ hence $\zeta \in S \Rightarrow \delta_\zeta \in \theta \backslash B \Rightarrow \delta_\zeta \notin B$; but $\delta_\zeta \in B$ by the definition of B, g^*, g_ζ, S.

Clause (a):

If $\epsilon \in \kappa \backslash w$, $\mathrm{cf}(\beta^*_\epsilon) > \theta$ then $\gamma^*_\epsilon =: \sup\{f_\alpha(\epsilon) : \alpha < \theta, f_\alpha(\epsilon) < \beta^*_\epsilon\}$ is $< \beta^*_\epsilon$ and it belongs to $N_{i(*)}$ (as $\epsilon, \langle f_\alpha : \alpha < \lambda \rangle$ and β^*_ϵ belongs to $N_{i(*)}$) and for any $\zeta \in S$ we get a contradiction to to $g_\zeta(\epsilon) = \beta^*_\zeta$.

Clearly $\zeta \in \kappa \backslash w \Rightarrow \ \mathrm{cf}[g^*(\zeta)] \geq \sigma$ (otherwise $g^*(\zeta) = \sup(N_i \cap g^*(\zeta))$ (as $N_i \prec (\mathscr{H}(\chi), \in, <^*_\chi)$ and $N_{i(*)} \cap \sigma = i(*)$ because $i(*) \in E$) and easy contradiction.

Clause (c):

If there is $\bar{\beta}' = \langle \beta'_\epsilon \in u \rangle$ contradicting clause (c), then there is such a sequence defined from $B, \langle f_\alpha : \alpha < \theta \rangle, d, w, \langle \beta^*_i : i < \kappa \rangle$, just use the $<^*_\chi$-first one, hence without loss of generality $\bar{\beta}' \in N_{i(*)}$, so for any $\zeta \in S$ we get a contradiction. $\square_{6.1}$

6.3 Observation. If $|\mathfrak{a}| < \min(\mathfrak{a}), H \subseteq \Pi\mathfrak{a}, |H| = \theta = \ \mathrm{cf}(\theta) \notin \ \mathrm{pcf}(\mathfrak{a})$ and also $\theta > \sup(\theta \cap \ \mathrm{pcf}(\mathfrak{a}))$ then for some $g \in \Pi\mathfrak{a}$, the set $H_g =: \{f \in H : g < g\}$ has cardinality θ; in face H is the union of $\leq \sup(\theta \cap \ \mathrm{pcf}(\mathfrak{a}))$ sets of the form H_g.
[Why? This is as $\Pi\mathfrak{a}/J_{<\theta}[\mathfrak{a}]$ is $\min(\mathrm{pcf}\mathfrak{a}) \backslash \theta)$-directed and the ideal $J_{<\theta}[\mathfrak{a}]$ is generated by $< \theta$ sets.

In details, let $\langle \mathfrak{b}_\sigma[\mathfrak{a}] : \sigma \in \ \mathrm{pcf}(\mathfrak{a}) \rangle$ be a generating sequence for $\mathfrak{a}$ (exists by [Sh:g, Ch.VIII], 2.6).

For $\sigma \in \ \mathrm{pcf}(\mathfrak{a})$ let $f^\sigma_\alpha \in \Pi(\mathfrak{a})$ for $\alpha < \sigma$ be such that $\langle f^\sigma_\alpha : \alpha < \sigma \rangle$ is $<_{J_\theta[\mathfrak{a}]}$-increasing and cofinal and moreover $\{f^\sigma_\alpha \restriction \mathfrak{b}_\sigma[\mathfrak{a}] : \alpha < \sigma\}$ is cofinal in $\Pi\mathfrak{b}_\sigma[\mathfrak{a}]$ (where $J_\sigma[\mathfrak{a}] = J_{<\sigma}[\mathfrak{a}] + \mathfrak{b}_\sigma[\mathfrak{a}]$), exists as $(\Pi\mathfrak{b}_\sigma, [\mathfrak{a}], <)$ has cofinality σ by [Sh:g, Ch.II], 3.1.

Now as $\Pi\mathfrak{a}/J_{<\theta}[\mathfrak{a}]$ is $\min(\mathrm{pcf}(\mathfrak{a}) \backslash \theta^+)$-directed (as $J_{<\theta}[\mathfrak{a}] = J_{<\theta^+}[\mathfrak{a}] = J_{<\min(\mathrm{pcf}(\mathfrak{a}) \backslash \theta^+)}[\mathfrak{a}]$ and by [Sh:g, Ch.I], 1.5) there is $g \in \Pi\mathfrak{a}$ such that: $h \in H \Rightarrow h > g \mod J_{<\theta}[\mathfrak{a}]$; hence for each $h \in F$ for some finite $\Theta(h) \subseteq \theta \cap \ \mathrm{pcf}(\mathfrak{a})$ we have

$$\{\sigma \in \mathfrak{a} : h(\sigma) \geq g(\sigma)\} \subseteq \bigcup\{\mathfrak{b}_\sigma[\mathfrak{a}] : \sigma \in \Theta(h)\}.$$

Also for every $\sigma \in \ \mathrm{pcf}(\mathfrak{a})$ we can find $\alpha_\sigma(h)$ (for $\sigma \in \ \mathrm{pcf}(\mathfrak{a}), h \in H$) such that $h \restriction \mathfrak{b}_\sigma[\mathfrak{a}] < f^\sigma_{\alpha_\sigma(h)} \restriction \mathfrak{b}_\sigma[\mathfrak{a}]$. So $h < \max(\{g, f^\sigma_{\alpha_\sigma(h)} : \sigma \in \Theta(h)\})$. Let

$$\begin{aligned} G =: \big\{\max\{g, f^{\sigma_1}_{\alpha_1}, \dots, f^{\sigma_n}_{\alpha_n}\} : & n < \omega, \{\sigma_1, \dots, \sigma_n\} \subseteq \theta \cap \ \mathrm{pcf}(\mathfrak{a}) \\ & \text{and } \alpha_1 < \sigma_1, \dots, \alpha_n < \sigma_n\big\} \end{aligned}$$

it has cardinality $\leq \aleph_0 + \sup(\theta \cap \text{ pcf}(\mathfrak{a}) < \theta$ and $\theta = \text{ cf}(\theta) = |H|$ and $\forall h \in H \exists g' \in G(h < g)$.
So for some $g^* \in G$ the set $\{h \in H : h < g^*\}$ has cardinality θ as required.] $\square_{6.3}$

We comment to [Sh 410], 3.7 (which solve a problem from Gerlits Hajnal Szentmiklossy [GHS]).

6.4 Claim. *1) Suppose γ^* and $i^* = i(*)$ are ordinals and $\bar{\chi} = \langle \chi_i : i < i^* \rangle$ is a sequence of infinite cardinals; so of course we can find $n, \bar{w} = \langle w_\ell : \ell < n \rangle, \bar{\kappa} = \langle \kappa_\ell : \ell < n \rangle, \bar{\sigma} = \langle \sigma_\ell : \ell < n \rangle$ such that $\bar{w}$ is a partition of $i^*, |w_\ell| = \kappa_\ell, \bigwedge\limits_{i \in w_\ell} \chi_i \leq \sigma_\ell$, and*

$$(\forall \chi < \sigma_\ell)(\exists^{\kappa_\ell} i)(i \in w_\ell \ \ \& \ \ \chi < \chi_i \leq \sigma_\ell)$$

and $\bar{\kappa}$ is strictly increasing, $\bar{\sigma}$ is strictly decreasing, in fact $\bar{w}, \bar{\kappa}, \bar{\sigma}$ are unique. Then the following are equivalent

$(A)_{\bar{\chi},\gamma^*}$ *there are $f_\alpha \in \prod\limits_{i < i(*)} \chi_i$ for $\alpha < \gamma^*$ satisfying $\alpha < \beta \Rightarrow (\exists i < i^*)[f_\alpha(i) > f_\beta(i)]$*

$(B)_{\bar{\chi},\gamma^*}$ *for some $\ell < n$ we have: $2^{\kappa_\ell} \geq |\gamma^*|$ or for every regular $\mu_1 \in (\gamma^* + 1) \backslash \sigma_\ell^+$ for some singular $\lambda^* \leq \sigma_\ell$ we have*

$(*)$ *$cf(\lambda^*) \leq \kappa_\ell, \lambda^* > 2^{\kappa_\ell}$ and $pp^+(\lambda^*) > \mu_1$*

2) If

$\otimes$ $(\forall \mu_1)(\mu_1 = \ cf(\mu_1) \leq |\gamma^*| \rightarrow (\exists \mu_2)[\mu_2 = \ cf(\mu_2) \ \ \& \ \ \mu_1 \leq \mu_2 \leq |\gamma^*| \ \ \& \ (\forall \alpha < \mu_2)|\alpha|^{\aleph_0} < \mu_2])$,

then in part (1), $(B)_{\bar{\chi},\gamma^*}$ the demand $(*)$ on λ^* can be replaced by

$(*)'$ *$cf(\lambda^*) \leq \kappa_\ell, \lambda^* > 2^{\kappa_\ell}$ and $(\forall \mu < \lambda^*)(\mu^{\kappa_\ell} < \lambda^*)$.*

Now we call it $(B)'_{\bar{\chi},\gamma^*}$ (so if $\otimes$ then $(A)_{\bar{\chi},\gamma^*} \Leftrightarrow (B)_{\bar{\chi},\gamma^*} \Leftrightarrow (B)'_{\bar{\chi},\gamma^*}$).
3) If $\otimes$ from part (2) holds, then also $(A)_{\bar{\chi},\gamma^*} \Leftrightarrow (B)''_{\bar{\chi},\gamma^*} \Leftrightarrow (B)^+_{\bar{\chi},\gamma^*}$ where

$(B)''_{\bar{c}hi,\gamma^*}$ $|\gamma^*| \leq \max\limits_{\ell < n} (\sigma_\ell)^{\kappa_\ell}$

$(B)^*$ *for some $\ell < n$ we have: $2^{\kappa_\ell} \geq |\gamma^*|$ or for every regular $\mu_1 \in (\gamma^* + 1) \backslash \sigma_\ell^+$ for some singular λ^* we have*

$(*)^+$ *$cf(\lambda^*) \leq \kappa_\ell, \lambda^* > 2^{\kappa_\ell}$ and $(\forall \mu < \lambda^*)(\mu^{\kappa_\ell} < \lambda^*)$ and $pp^+(\lambda^*) > \mu_1$.*

4) In part (2), (3) instead $\otimes$ we may let $\lambda_0 = \max\{2^{\kappa_\ell} : \ell < n\}, \lambda_1 = |\gamma^*|$, demand

$\oplus_{\lambda_0,\lambda_1}$ *if $\lambda_0 < \mu \leq \lambda_1$, $cf(\mu) = \aleph_0$ and $(\forall \lambda < \mu)(|\lambda|^{\aleph_0} < \mu)$ then $pp(\mu) =^+ \mu^{\aleph_0}$.*

Remark. 1) On $\oplus_{\lambda_0,\lambda_1}$ from 6.4(4), see [Sh 430],§1.
2) Note that we could in 6.4(1) demand

$$(\forall \chi < \sigma_\ell)(\exists^{\kappa_\ell} i)(i \in {}^w\ell \ \ \& \ \ \chi \le \chi_1 < \sigma_\ell)$$

and can allow χ_i (hence σ_i) to be any ordinal, and even let κ_i be ordinal so the demand is $(\forall \alpha < \sigma_\ell)[\kappa_\ell = \text{ otp}\{i \in w_\ell : \alpha \le \chi_i < \sigma_\ell\}]$. This causes no serious change.

Proof. 1) $(B)_{\bar{\chi},\gamma^*} \Rightarrow (A)_{\bar{\chi},\gamma^*}$.

Let $\ell < n$ exemplifies $(B)_{\bar{\chi},\gamma^*}$ so there are $f'_\alpha \in {}^{\kappa_\ell)}(\sigma_\ell)$ for $\alpha < \gamma^*$ such that $\alpha < \beta < \gamma^* \Rightarrow (\exists j < \kappa_\ell)[f'_\alpha(j) > g'_\beta(j)]$.
[Why? Easy by cases.]

Let $h : w_\ell \rightarrow \kappa_\ell$ be such that:

$$(\forall j < \kappa_\ell)(\forall \sigma < \sigma_\ell)(\exists^{\kappa_\ell} i)[i \in w_\ell \ \ \& \ \ \sigma < \chi_i \le \sigma_\ell \ \ \& \ \ h(i) = j].$$

Let $f_\alpha \in \prod\limits_{i<i(*)} \chi_i$ be: $f_\alpha(i) = f'_\alpha(h(i))$ if $i \in w_\ell, f'_\alpha(h(i)) < \chi_i$ and $f_\alpha(i) = 0$ otherwise; so if $\alpha < \beta < \gamma^*$ then for some $j < \kappa_\ell, f'_\alpha(j) < f'_\beta(j) < \sigma_\ell$ so for some $i \in w_\ell$ we have: $h(i) = j \ \ \& \ \ \chi_i > f'_\beta(j)$. So $f_\alpha(i) = f'_\alpha(j) < f'_\beta(j) < f_\beta(i)$ as required.

$(A)_{\bar{\chi},\gamma^*} \Rightarrow (B)_{\bar{\chi},\gamma^*}$

Assume this fails, note that clearly $\gamma_1 < \gamma_2 \ \ \& \ \ (A)_{\bar{\chi},\gamma_2} \Rightarrow (A)_{\bar{\chi},\gamma_1}$ and $\gamma_1 < \gamma_2 \ \ \& \ \ (B)_{\bar{\chi},\gamma_2} \Rightarrow (B)_{\bar{\chi},\gamma_1}$.Without loss of generality γ^* is minimal (for our $\bar{c}hi$) for which this fails; so γ^* is minimal such that $(B)_{\bar{\chi},\gamma^*}$ fails inspecting $(B)_{\bar{\chi},\gamma^*}$ as n is finite, clearly γ^* is a regular cardinal, call it θ. By renaming we can assume that i^* is a cardinal. Now let $\langle f_\alpha : \alpha < \theta\rangle$ exemplifies $(A)_{\bar{\chi},\theta}$; now apply 6.1 above with $i^*, \langle f_\alpha : \alpha < \theta\rangle, D^{cb}_\theta$ the filter of cobounded subsets of $\theta, \max\limits_{\ell<n}(2^{\kappa_\ell})^+ = (2^{|i^*|})^+$ here standing for $\kappa, \langle\langle\beta^\alpha_i : i < \kappa\rangle : \alpha < \theta\rangle, D, \sigma$ there.

So we get $w, \langle\beta^*_i : i < i^*\rangle, B$ as there, and let $\mathfrak{a} = \{\text{cf}(\beta^*_i) : i \in i^*\backslash w\}$, so $\mathfrak{a} \subseteq \text{ Reg } \cap (\max\limits_{\ell<n}\sigma_\ell)\backslash(2^{|i^*|})^+$ (see clause (a) of 6.1) and $|\mathfrak{a}| \le |i^*|$. Now if $\theta \le \max \text{pcf}(\mathfrak{a})$ then for some $\ell, \theta \le \max \text{pcf}\{\text{cf}(\beta^*_i) : i \in w_\ell\backslash w\}$, and so $(B)_{\bar{\chi},\theta}$ does not fail, contradicting an earlier assumption. So $\theta > \max \text{pcf}(\mathfrak{a})$, so there is a cofinal $H \subseteq \prod\limits_{i\in(i^*\backslash w)} \beta^*_i$ of cofinality $< \theta$, so there are $h_\alpha \in H$ such that $f_\alpha \restriction (i^*\backslash w) < h_\alpha$ but $|B| = \theta > |H|$ (by the choice of D^{cb}_θ as D) so for some $h^* \in H$ the set $B_1 = \{\alpha \in B : h_\alpha = h^*\}$ is unbounded in θ. By clause (c) of the conclusion of 6.1 for some $\alpha \in B$ we have $i \in i^*\backslash w \Rightarrow h^*(i) < f_\alpha(i)$. Choose $\beta \in B_1\backslash(\alpha + 1)$, so $\alpha < \beta$ are in B, hence $f_\alpha \restriction w = f_\beta \restriction w$ and $i \in i^*\backslash w \Rightarrow f_\beta(i) < h^*(i) < f_\alpha(i)$, so $\bigwedge\limits_i f_\beta(i) \le f_\alpha(i)$, a contradiction to $\langle f_\gamma : \gamma < \theta\rangle$ exemplifies $(A)_{\bar{\chi},\theta}$.

2), 3) Clearly $(B)^+_{\bar{\chi},\gamma^*} \Rightarrow (B)_{\bar{\chi},\gamma^*} \Rightarrow (B)''_{\bar{\chi},\gamma^*}$ by checking. So we should just prove:

$(B)''_{\bar{\chi},\gamma^*} \Rightarrow (B)'_{\bar{\chi},\gamma^*}$

We can assume $|\gamma^*| > 2^{\kappa_\ell}$ for $\ell < n$ (otherwise the conclusion is trivial). We know by $(B)''_{\bar{\chi},\gamma^*}$ that for some $\ell, (\sigma_\ell)^{\kappa_\ell} \ge |\gamma^*|$. Let us check now $(B)'_{\bar{\chi},\gamma^*}$ so let a regular

$\mu_1 \in (\gamma^* + 1) \backslash \sigma^+_\ell$ be given. So $(\sigma_\ell)^{\kappa_\ell} \geq |\gamma^*| \geq \mu_1$ hence $\lambda =: \min\{\lambda : \lambda^{\kappa_\ell} \geq \mu_1\}$ is $\leq \sigma_\ell$ but is $> 2^{\kappa_\ell}$, hence it is singular, $\mathrm{cf}(\lambda) \leq \kappa_\ell$ and $(\forall \alpha < \lambda)(|\alpha|^{\kappa_\ell} < \lambda)$, i.e. as required in $(*)'$.

$(B)'_{\bar{\chi},\gamma^*} \Rightarrow (B)^+_{\bar{\chi},\gamma^*}$

Again we can assume $|\gamma^*| > 2^{\kappa_\ell}$ for $\ell < n$. Let us check $(B)_{\bar{\chi},\gamma^*}$, so let a regular $\mu_1 \in (\gamma^* + 1) \backslash \sigma^+_\ell$ be given. As we are assuming $\otimes$ from 6.4(2) there is $\mu_2 \in$ Reg $\cap \, (\gamma^* + 1) \backslash \mu_1$ such that $(\forall \alpha < \mu_2)(|\alpha|^{\aleph_0} < \mu_2)$. Apply $(B)'_{\chi,\gamma^*}$ for μ_2 and get λ^*_2 as in $(*)$ for μ_2 instead of μ_1. Clearly $(\lambda^*_2)^{\kappa_\ell} \geq \mu_2$ and let $\lambda^* = \min\{\lambda : \lambda^{\kappa_\ell} \geq \mu_2\}$, so $\lambda^* \leq \lambda^*_2 \leq \sigma_\ell, \lambda^* > 2^{\kappa_\ell}$ and $(\forall \mu < \lambda^*)[\mu^{\kappa_\ell} < \lambda^*]$, and clearly $\mathrm{cf}(\lambda^*) \leq \kappa_\ell$, so $(\lambda^*)^{\kappa_\ell} = (\lambda^*)^{\mathrm{cf}(\lambda)}$.

By the choice of μ_2 necessarily $\mathrm{cf}(\lambda^*) > \aleph_0$ (otherwise $\mu_2 \leq (\lambda^*)^{\kappa_\ell} = (\lambda^*)^{\aleph_0} < \mu_2$). By [Sh:g], (see 6.5 below), $\mathrm{pp}(\lambda^*) =^+ (\lambda^*)^{\mathrm{cf}(\lambda^*)} = (\lambda^*)^\kappa$ as required in $(*)^+$.
4) The only place we use was in choosing μ_2 in the proof of $(B)'_{\bar{\chi},\gamma^*} \Rightarrow (B)^+_{\bar{\chi},\gamma^*}$ and the use of its property is to show $\mathrm{cf}(\lambda^*) > \aleph_0$ (to be able to use [Sh:g, Ch.VIII],§1) but we can use instead $\oplus_{\lambda_0,\lambda_1}$. $\square_{6.4}$

Remember that by [Sh:g]:

6.5 Observation. If μ is singular, $\mathrm{cf}(\mu) > \aleph_0$ and $\alpha < \mu \Rightarrow |\alpha|^{\mathrm{cf}(\mu)} < \mu$ then $\mu^{\mathrm{cf}(\mu)} = \mathrm{pp}(\mu)$.

Proof. By [Sh 430], 3.5, [Sh:g, CH.VIII], 1.8, [Sh:g, Ch.II], **?**. $\square_{6.5}$

—> scite{5.6} undefined

6.6 Definition. 1) For $F \subseteq {}^\delta\mathrm{Ord}$, we say F is free$^\ell$ when we can find $\zeta_f < \delta$ for $f \in F$ such that:

(a) if $\ell = 1$ then

$$f \neq g \in F, \zeta = \max\{\zeta_f, \zeta_g\} \Rightarrow f \restriction \zeta \neq g \restriction \zeta$$

(b) if $\ell = 2$ then

$$f \neq g \in F, \delta > \zeta \geq \max\{\zeta_f, \zeta_g\} \Rightarrow f(\zeta) \neq g(\zeta)$$

(c) if $\ell = 3$ then

$$f \neq g \in F, \delta > \zeta > \varepsilon = \max\{\zeta_f, \zeta_g\}, f(\varepsilon) \leq g(\varepsilon) \Rightarrow f(\zeta) \leq g(\zeta)$$

(d) if $\ell = 4$ then for $f, g \in F, f \leq g$ (i.e.
$\zeta < \delta \Rightarrow f(\zeta) \leq g(\zeta)$) or $g \leq f$

(e) if $\ell = 5$ then for some ζ and h we have

$$f \in F \Rightarrow f \restriction \zeta = h$$

$$\{f(\zeta) : f \in F\} \text{ is with no repetition.}$$

2) Let free$^{\ell,m}$ means free$^\ell$ and free^m. For J an ideal on δ we write J-free$^\ell$ if $\zeta_f < \delta$ is replaced by $s_f \in J$.

6.7 Definition. For $F \subseteq {}^{\delta}\mathrm{Ord}$:
1) We say F is μ-free^x if every $F' \in [F]^{\mu}$ is free^x.
2) We say F is (μ, κ)-free^x if every $F' \in [F]^{\mu}$ there is $F'' \in [F']^{\kappa}$ which is free^x.

6.8 Fact: 1) "F is free$^{\ell}$ implies F is free^{m}" when (ℓ, m) is one of (2,1), (4,3), (5,1).
2) Similarly for μ-free^x and (μ, κ)-free^x.

Proof. Straight.

On 6.9 see [Sh:g, Ch.II],§1, [Sh:g, Ch.II],§3, [Sh 282], [Sh:g, Ch.II], 4.10, Shelah Zapletal [ShZa 561].

6.9 Claim. *1) If* $J^{bd}_{\delta} \subseteq J, J$ *an ideal on* $\delta, \prod\limits_{i<\delta} \lambda_i / J$ *is* λ^+*-directed,* $\langle \lambda_i : i < \delta \rangle$ *an increasing sequence of regulars* $> \delta$ *with limit* $\mu, \mu < \lambda = cf(\lambda)$, *then there are regulars* $\lambda'_i < \lambda_i$ *with* $tlim(\lambda'_i) = \mu$ *and* $f_\alpha \in \Pi \lambda'_i$ *for* $\alpha < \lambda$ *such that* $\langle f_\alpha : \alpha < \lambda \rangle$ *is* $<_J$*-increasing cofinal in* $\prod\limits_{i<\delta} \lambda'_i$ *and* $\{f_\alpha : \alpha < \lambda\}$ *is* $\mu^+ - J$*-free.*
2) Assume $\mathfrak{a}$ *is a set of regular* $> |\mathfrak{a}|$ *with no last element,* J *an ideal on* $\mathfrak{a}$ *extending* $J^{bd}_{\mathfrak{a}}$ *and* $\mathfrak{c} = \{\theta \in \mathfrak{a} : \theta > \max pcf(\theta \cap \mathfrak{a})\}$ *and* $\lambda = \max pcf(\mathfrak{a})$. *Then there is* $\langle f_\alpha : \alpha < \lambda \rangle$ *cofinal in* $\Pi \mathfrak{a}, <_J$*-increasing, such that:*

$(*)$ *if* $\theta \in \mathfrak{c}$ *then* $\{f_\alpha \restriction (\theta \cap \mathfrak{a}) : \alpha < \lambda\}$ *has cardinality* $< \theta$.

3) If $\mu > \mu_0 \geq \kappa \geq cf(\mu), \lambda = \mu^+$ *(or just* $\mu < \lambda = cf(\lambda) < pp^+_\kappa(\mu)$ *then for some* $\mathfrak{a} \subseteq (\mu_0, \mu) \cap Reg, |\mathfrak{a}| \leq \kappa, [\theta \in \mathfrak{a} \Rightarrow \max pcf(\theta \cap \mathfrak{a}) < \theta]$ *and* $\lambda = \max pcf(\mathfrak{a})$ *(if* $[\alpha < \mu \Rightarrow |\alpha|^\kappa < \mu]$ *we can have* $\mu = \sup(\mathfrak{a})$, $otp(\mathfrak{a}) = cf(\mu)$ *(so part (2) is not empty)).*
4) If J *is an ideal on* $\mathfrak{a}, \langle f_\alpha : \alpha < \lambda \rangle$ *is* $<_J$*-increasing* $<_J$*-cofinal in* $\Pi \mathfrak{a}$ *and* J *is generated by* $< \min(\mathfrak{a})$ *sets (as an ideal) then every* $A \in [\lambda]^\lambda$ *for some* $\mathfrak{d} \in J$ *we have: for every* $g \in \Pi \mathfrak{a}$ *for* λ *ordinals* $\alpha \in A, g \restriction (\mathfrak{a} \backslash \mathfrak{d}) < f_\alpha \restriction (\mathfrak{a} \backslash \mathfrak{d})$. *Hence* $\bar{f}$ *is* $(\lambda, \min(\mathfrak{a}) - J$*-free* $^{\{2,3\}}$.
5) Assuming $|\mathfrak{a}| < \min(\mathfrak{a}), \lambda = tcf(\Pi \mathfrak{a}, <_{J^{bd}_{\mathfrak{a}}}), \mathfrak{c} = \{\theta \in \mathfrak{a} : \max pcf(\theta \cap \mathfrak{a})\}$ *is unbounded in* $\mathfrak{a}$ *and* $\langle f_\alpha : \alpha < \lambda \rangle$ *as in part 2. Then not only for each* $\theta \in \mathfrak{a}$ *is* $\{f_\alpha : \alpha < \lambda\}, (\lambda, \theta)$*-free*$^{\{2,3\}}$, *but for any* $A \in [\lambda]^\lambda$ *for every large enough* $\theta \in \mathfrak{c}$ *there is* $B \in [A]^\theta$ *such that* $\langle f_\alpha \restriction (\theta \cap \mathfrak{a}) : \alpha \in B \rangle$ *is constant and* $\langle f_\alpha \restriction (\mathfrak{a} \backslash \theta) : \alpha \in B \rangle$ *is strictly increasing.*
6) Assume $\lambda = tcf(\Pi \mathfrak{a}, <_J), J^{bd}_{\mathfrak{a}} \subseteq J, \lambda > \mu = \sup(\mathfrak{a})$ *and* $\langle f_\alpha : \alpha \in \lambda \rangle$ *is* $<_J$*-increasing and* $<_J$*-cofinal*

(a) *if* $\kappa \leq \mu^+$ *and*[3] $\{\delta < \lambda : cf(\delta) < \kappa\} \in I[\lambda]$, *then for some* $A \in [\lambda]^\lambda$ *we have* $\{f_\alpha : \alpha \in A\}$ *is* μ*-free*^{1}, *also we can find* $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$ *as above with* $A = \lambda$, *such that* $\bar{f}$ *is* ^{b} *continuous (see [Sh:g, Ch.I,§3])*

(b) *if* $\kappa = cf(\kappa) < \mu^+$ *and* $\{\delta < \lambda : cf(\delta) < \kappa\}$ *we get similar results for* (κ, κ)*-free*^{1}.

---

[3] note: if λ is a successor of regular $> \kappa^+$ then this holds

7) Assume $\langle \mu_i : i \le \kappa \rangle$ *is an increasing continuous sequence of singulars* $> \kappa, \kappa = cf(\kappa) > \aleph_0, \theta \in Reg \cap \mu_0 \backslash \kappa^+$ *and*[4] $\delta < \mu_i^+ : cf(\delta) = \theta\} \in I[\mu_i^+]$ *for* $i \le \kappa$ *and* $pp_\kappa(\mu_i) < \mu_{i+1}^+$ *for each* $i \le \kappa$. *Then* $<_{J^{bd}_C}$*-increasing and cofinal and is* (θ, θ)*-free*^{1}. *If we demand in addition* $\{\delta < \mu_i^+ : cf(\delta) \le \theta\} \in I[\mu_i^+]$ *then* F *is* θ^+*-free* ^{1}.

Proof. 1) By [Sh:g, Ch.II],§1.
2) By [Sh:g, Ch.II], 3.5.
3) By [Sh:g, Ch.II],§3.
4) Can prove as in [Sh 282]. Or as in 6.1, as below.

Let $A \in [\lambda]^\lambda$ and let $\{\mathfrak{d}_\zeta : \zeta < \zeta^*\} \subseteq J$ be a family generating J closed under finite union such that $\zeta^* < \lambda_0$. We shall prove that for some $\zeta < \zeta^*$ we have $(\forall g \in \Pi\mathfrak{a})(\exists^\lambda \alpha \in A)(g \restriction (\mathfrak{a} \backslash \mathfrak{d}_\zeta) < f_\alpha \restriction (\mathfrak{a} \backslash \mathfrak{d}_\zeta))$. If not, for each $\zeta < \zeta^*$ some $g_\zeta \in \Pi\mathfrak{a}$ and $\alpha_\zeta < \lambda$ exemplifies it, i.e. $\alpha \in A \backslash \alpha_\zeta \Rightarrow \neg(g \restriction (\mathfrak{a} \backslash \mathfrak{d}_\zeta) < f_\alpha \restriction (\mathfrak{a} \backslash \mathfrak{d}_\zeta))$. So defined by the function $g \in \Pi\mathfrak{a}, g(\theta) =: \sup[\{g_\zeta(\theta) : \zeta < \zeta^*\} \cup \{f_{\alpha^*}(\theta) + 1\}]$ is well defined and let $\alpha^* = \sup\{\alpha_\zeta : \zeta < \zeta^*\} < \lambda$. Now $\{f_\alpha : \alpha < \lambda\}$ is $<_J$-increasing and $<_J$-cofinal in $\Pi\mathfrak{a}$ so for some $\alpha \in A \backslash (\alpha^* + 1), g <_J f_\alpha$, so for some $\zeta < \zeta^*$ we have $\{\theta \in \mathfrak{a} : \neg(g(\theta) < f_\alpha(\theta))\} \subseteq \mathfrak{d}_\zeta$, so for $\mathfrak{d}_\zeta$, the pair (g_ζ, α_ζ) is not as required, a contradiction.
5), 6, 7) Left to the reader. $\square_{6.9}$

6.10 Definition. We say $(\Pi\mathfrak{a}, <_J)$ is X-free^y if there is a $<_J$-increasing $<_J$-cofinal $\langle f_\alpha : \alpha < \lambda \rangle$ from $\Pi\mathfrak{a}$ which is y-free^x.

6.11 Fact: If $y \in \{1, 2, 3, \{2, 3\}\}, x \in \{\mu - J, (\mu, \theta) - J\}$ and $(\Pi\mathfrak{a}, <_J)$ is x-free^y and $\langle f_\alpha : \alpha < \lambda \rangle$ is $<_J$-increasing $<_J$-cofinal in $\Pi\mathfrak{a}$ then for some $A \in [\lambda]^\lambda$ we have $\{f_\alpha : \alpha \in A\}$ is x-free^y.

Proof. Straight.

6.12 Question: Let $\mu > \kappa \ge$ cf(μ). For how many $\mathfrak{a} \subseteq$ Reg $\cap \mu, \mu = \sup(\mathfrak{a}), J^{bd}_{\mathfrak{a}} \subseteq J, J$ an ideal on $\mathfrak{a}, \lambda =$ tcf$(\Pi\mathfrak{a}, <_J)$, is $(\Pi\mathfrak{a}, <_J)$ not μ-free?

The proof of [Sh:g, Ch.IX], 3.5 has a gap (in the reference to [Sh:g, Ch.IX], **?**).
—> scite{3.3A} undefined
What we know is only

6.13 Lemma. *1) Assume* $cf(\lambda) \in [\sigma, \theta), \lambda > \theta > cf(\lambda) \ge \sigma = cf(\sigma) > \aleph_0$. *Then* $cov(\lambda, \lambda, \theta, \sigma) =^+ \sup\{pcf_{\Gamma(\theta,\sigma), J^{bd}_{\mathfrak{a}}}(\mathfrak{a}) : \mathfrak{a} \subseteq Reg \cap \lambda, \lambda = \sup(\mathfrak{a}), |\mathfrak{a}| < \min(\mathfrak{a})\}$ *where*

$$pcf_{\Gamma(\theta,\sigma),J}(\mathfrak{a}) = \big\{ tcf(\Pi\mathfrak{a}, <_I) : I \text{ an ideal on } \mathfrak{a} \text{ extending } J, tcf(\Pi\mathfrak{a}, <_I) \text{ well defined, } I \text{ is } \sigma\text{-complete and for some } \mathfrak{b} \in I, |\mathfrak{a} \backslash \mathfrak{b}| < \theta \big\}$$

---

[4] note: if λ is a successor of a regular $> \kappa^+$ then this holds

(the $=^+$ means that if the left side is regular then the supremum in the right side is obtained).
2) If in addition $(\forall \mu < \lambda)(cov(\mu,\theta,\theta,\sigma) < \lambda), (\theta, \sigma$ *regular)* *then*

$$cov(\lambda,\lambda,\theta,\sigma) =^+ \ pp_{\Gamma(\theta,\sigma)}(\lambda).$$

3) So for $\mathscr{Y} = \mathscr{Y}_\mu, Eq = Eq_\mu$ *be as in [Sh 420, §3- §5],* $cf(\mu) = \sigma > \aleph_0$ *for simplicity,* $\mu > \theta > \ cf(\mu)$*.*
If $\mathfrak{a} \subseteq \ Reg \setminus \mu, |\mathfrak{a}| < \mu, |\mathfrak{a}| < \min(\mathfrak{a}), \lambda$ *inaccessible,* $J = J^{bd}_{\mathfrak{a}}, \lambda = \ sup\ pcf_{\Gamma(\theta,\sigma),J^{bd}}(\mathfrak{a})$*, then we can find* $e \in Eq, \bar\lambda = \langle \lambda_x : x \in \mathscr{Y}_\mu/e\rangle$ *and* $D \in FIL(\mathscr{Y}_\mu)$ *such that:*

$$\lambda = \ tcf(\Pi\bar\lambda/D)$$
$$lim_D(\bar\lambda) = \mu$$
$$\lambda_x = \ cf(\lambda_x).$$

Proof. As in [Sh 410],§1 (replacing normal by σ-complete) or make the following changes in the proof of [Sh:g, Ch.IX], 3.5: $\|N^x_k\| = \mu_k, \mu_k + 1 \subseteq N^x_k$.

6.14 Claim. *1) Assume*

- (i) λ *is inaccessible*
- (ii) $|\mathfrak{a}|^+ < \min(\mathfrak{a}), \mu =: \sup(\mathfrak{a}) < \lambda$
- (iii) $R \subseteq \lambda \cap \ Reg \setminus \mu, |R| = \lambda,$
- (iv) *for* $\tau \in R, \mathfrak{b}_\tau \subseteq \mathfrak{a}, \sup(\mathfrak{b}_\tau) = \mu, J_\tau$ *an ideal on* $\mathfrak{b}_\theta$ *including* $J^{bd}_{\mathfrak{b}_\theta}, \tau = tcf(\Pi\mathfrak{b}_\theta, <_{J\theta})$.

Then for some $\langle \lambda_\theta : \theta \in \mathfrak{a}\rangle$ *we have*

- (a) $\lambda_\theta = \ cf(\lambda_\theta) < \theta, \lambda_\theta > |\mathfrak{a}|$
- (b) $lim_{J_\tau}\lambda_\theta = \mu$
- (c) $\lambda = \ tcf(\prod\limits_{\theta\in\mathfrak{a}} \lambda_\theta, <_{J_{<\lambda}[\mathfrak{a}]})$
- (d) $\{\min\ pcf_{J_\tau}(\prod\limits_{\theta\in\mathfrak{b}_\tau} \lambda_\theta, <_{J_\tau}) : \tau \in R\}$ *is unbounded in* λ.

So $\mathfrak{a}' =: \{\lambda_\theta : \theta \in \mathfrak{a}\}$ *satisfies (ii), (iii), (iv) and*

- $(i)^-$ $|\mathfrak{a}'|^+ < \min(\mathfrak{a}')$ *instead*
- (i) $|\mathfrak{a}|^+ < min(\mathfrak{a})$*, immaterial and*
- (v) *max* $pcf(\mathfrak{a}') = \lambda$.

2) Assume $(i)^-, (ii), (iii), (iv), (v)$ *are satisfied by* $\mathfrak{a}, R, \mathfrak{b}_\tau, J_\tau$*. Then for some* $f_\tau \in \Pi\mathfrak{b}_\tau$ *for* $\tau \in R$ *we have:*

- $(*)$ *for every* $g \in \Pi\mathfrak{a}$ *for some* $\tau, g \restriction \mathfrak{b}_\tau <_{J_\tau} f_\tau$.

Proof. 1) By [Sh:g, Ch.II], **?** we can find $\langle \lambda_\theta : \theta \in \mathfrak{a} \rangle$ satisfying (a), (b), (c). If
$\longrightarrow$ scite{1.5A} undefined
(d) fails choose for each $\tau \in R, \mathfrak{b}'_\tau \in J^+_\tau$ such that $\chi_\tau = \text{ tcf}(\prod\limits_{\theta \in \mathfrak{b}'_\tau} \lambda_\theta, <_{J_{\tau \restriction \mathfrak{b}'_\tau}})$ is well defined and equal to min $\text{pcf}_{J_\tau}(\prod\limits_{\theta \in \mathfrak{b}_\tau} \lambda_\theta, <_{J_\tau})$. So $\{\chi_\tau : \tau \in R\}$ is unbounded in λ hence for some $\chi, R' = \{\tau \in R : \chi_\tau = \chi\}$ is unbounded in λ. Hence we can find $\zeta^* < |\mathfrak{a}|^+$ and $\tau_\zeta \in R'$ for $\zeta < \zeta^*$ such that $\lambda \le \text{ max pcf}\{\tau_\zeta : \zeta < \zeta^*\}$ (choose by induction on $\zeta < |\mathfrak{a}|^+, \tau_\zeta \in R', \tau_\zeta > \text{ max pcf}\{\tau_\varepsilon : \varepsilon < \zeta\}$ and use localization). Let D_ζ be an ultrafilter on $\mathfrak{a}, b'_{\tau_\zeta} \in D_\tau, J_{\tau_\zeta} \cap D = \emptyset$, so $\text{tcf}(\Pi \mathfrak{a}, <_{D_\zeta}) = \tau_\zeta$, and E is an ultrafilter on $\{\tau_\zeta : \zeta < \zeta^*\}$ such that $\text{tcf}(\prod\limits_\zeta \tau_\zeta, <_E) \ge \lambda$. Let $D = \{\mathfrak{c} \subseteq \mathfrak{a} : \{\zeta : \mathfrak{c} \in D_\zeta\} \in E\}$, so $\text{tcf}(\Pi \mathfrak{a}, <_D) \ge \lambda$ hence $D \cap J_{<\lambda}[\mathfrak{a}] = \emptyset$. Apply this to $\langle \lambda_\theta : \theta \in \mathfrak{a} \rangle$, by [Sh:g, Ch.I], **? - ?** we get a contradiction.
$\longrightarrow$ scite{1.10} undefined
$\longrightarrow$ scite{1.11} undefined
3) Let $\langle f^*_\alpha : \alpha < \lambda \rangle$ be $<_{J_{<\lambda}[\mathfrak{a}]}$-increasing and cofinal. Let $f_\alpha = f^*_\alpha \restriction \mathfrak{b}_\alpha$. $\square_{6.14}$

Concerning [Sh 430],3.1 we comment

6.15 Theorem. *1) Assume $\lambda > \theta > \aleph_0$ are regular and*

$(*)_{\theta,\kappa}$ *if $\mathfrak{a} \subseteq$ Reg $\cap \lambda \backslash \theta$ and $|\mathfrak{a}| < \theta$ then there are $\zeta^* < \theta$ and $\mathfrak{b}_\zeta \in J_{<\lambda}[\mathfrak{a}]$ for $\zeta < \zeta^*$ such that for every $\mathfrak{b} \subseteq \mathfrak{a}, |\mathfrak{b}| < \kappa$ for some $\zeta < \zeta^*, \mathfrak{b} \subseteq \mathfrak{b}_\zeta$.*

Then the following conditions are equivalent:

$(A) = (A)_{\lambda,\theta,\kappa}$ *for every $\mu < \lambda$ we have cov$(\mu, \theta, \kappa, 2) < \lambda$*

$(B) = (B)_{\lambda,\theta,\kappa}$ *if $\mu < \lambda$ and $a_\alpha \in \mathscr{S}_{<\kappa}(\mu)$ for $\alpha < \lambda$ then for some $W \subseteq \lambda$ of cardinality λ we have $|\bigcup\limits_{\alpha \in W} a_\alpha| < \theta$*

$(C) = (C)_{\lambda,\theta,\kappa}$ *if a_α is a set of cardinality $< \kappa$ for $\alpha < \lambda$ and $W_0 \subseteq \{\delta < \lambda : cf(\delta) \ge \kappa\}$ then for some stationary $W \subseteq W_0$ and set b of cardinality $< \theta$ we have $\langle a_\alpha \backslash b : \alpha \in W \rangle$ is a sequence of pairwise disjoint sets.*

2) If $\lambda > \theta_1 > \theta_2 \ge \kappa > \aleph_0$ where λ, κ are regular, $(A)_{\lambda,\theta,\kappa} \Leftrightarrow (B)_{\lambda,\theta_1,\kappa}$ and cov$(\theta_1, \theta_2, \kappa, 2) < \lambda$ then $(A)_{\lambda,\theta_2,\kappa} \Leftrightarrow (B)_{\lambda,\theta_2,\kappa}$ (so if for some $\theta_1, (*)_{\theta_1,\kappa}, \lambda > \theta_1 > \theta_2 =$ cf$(\theta) \ge \kappa$ and cov$(\theta_1, \theta, \kappa, 2) < \lambda$ then the conclusions holds).

Proof. 1) Read the proof of [Sh 430], 3.1 (which was written in a way appropriate to this generalization), but defining the $M_n, \langle N^n_\zeta : \zeta < \theta \rangle$, we omit clause (d), that is, $N^n_\zeta \in \mathfrak{A}_{\delta(*)}$ and instead demand

$(d)'$ for each n we can find $\mathscr{P}_n \subseteq [\theta]^{<\theta}$ such that $(\forall a \in [\theta]^{<\kappa})(\exists b \in \mathscr{P})(a \subseteq b)$ and $\langle N^n_b : b \in \mathscr{P}_n \rangle$ such that $N^n_b \prec \mathfrak{B}, N^n_\zeta = \bigcup \{N^n_b : b \in \mathscr{P}_n, b \subseteq \zeta\}$ and $b_1 \subseteq b_2 \Rightarrow N^n_{b_1} \prec N^n_{b_2}$.

2) Left to the reader.

6.16 Question: Can we in [Sh 430], 4.2(1) weaken clause (β) in the conclusion to "$\lambda_x > \mu_0$ for D-almost all $x \in \mathscr{Y}/e$" then we can weaken the hypothesis [Sh 420], **?** (was stated in [Sh 430], earlier version clear).

—> scite{6.1C} undefined

REFERENCES.

[FShS 544] Sakaé Fuchino, Saharon Shelah, and Lajos Soukup. Sticks and clubs. *Annals of Pure and Applied Logic*, **to appear**.

[FShS 543] Sakaé Fuchino, Saharon Shelah, and Lajos Soukup. On a theorem of Shapiro. *Mathematica Japonica*, **40**:199–206, 1994.

[GHS] J. Gerlits, Andras Hajnal, and Z. Szentmiklossy. On the cardinality of certain Hausdorff spaces. *Discrete Mathematics*, **108**:31–35, 1992. Topological, algebraical and combinatorial structures. Frolik's Memorial Volume.

[Sh:E11] S. Shelah. An answer to complaints of E.Weitz on [Sh:371] and other complaints on [Sh:g].

[Sh 589] Saharon Shelah. PCF theory: applications. *Journal of Symbolic Logic*, **submitted**.

[Sh 620] Saharon Shelah. Special Subsets of ${}^{\mathrm{cf}(\mu)}\mu$, Boolean Algebras and Maharam measure Algebras. *General Topology and its Applications - Proc. of Prague Topological Symposium 1996*, **accepted**.

[Sh 460] Saharon Shelah. The Generalized Continuum Hypothesis revisited. *Israel Journal of Mathematics*, **accepted**.

[Sh 282] Saharon Shelah. Successors of singulars, cofinalities of reduced products of cardinals and productivity of chain conditions. *Israel Journal of Mathematics*, **62**:213–256, 1988.

[Sh 420] Saharon Shelah. Advances in Cardinal Arithmetic. In *Finite and Infinite Combinatorics in Sets and Logic*, pages 355–383. Kluwer Academic Publishers, 1993. N.W. Sauer et al (eds.).

[Sh 410] Saharon Shelah. More on Cardinal Arithmetic. *Archive for Mathematical Logic*, **32**:399–428, 1993.

[Sh 371] Saharon Shelah. Advanced: cofinalities of small reduced products. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter VIII. Oxford University Press, 1994.

[Sh 355] Saharon Shelah. $\aleph_{\omega+1}$ has a Jonsson Algebra. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter II. Oxford University Press, 1994.

[Sh:g] Saharon Shelah. *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.

[Sh 400] Saharon Shelah. Cardinal Arithmetic. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter IX. Oxford University Press, 1994. Note: See also [Sh400a] below.

[Sh 430] Saharon Shelah. Further cardinal arithmetic. *Israel Journal of Mathematics*, **95**:61–114, 1996.

[ShZa 561] Saharon Shelah and Jindra Zapletal. Embeddings of Cohen algebras. *Advances in Mathematics*, **126**(2):93–118, 1997.